

Poster Sessions – Abstract P002

Factors associated with the continuum of care of HIV-infected patients in Belgium

Van Beckhoven, Dominique¹; Lacor, Patrick²; Moutschen, Michel³; Piérard, Denis⁴; Sasse, André¹; Vaira, Dolorès⁵; Van den Wijngaert, Sigi⁶; Vandercam, Bernard⁷; Van Ranst, Marc⁸; Van Wijngaerden, Eric⁹; Vandekerckhove, Lino¹⁰; Verhofstede, Chris¹¹; Verbrugge, Ruth¹; Demeester, Rémy¹²; De Wit, Stéphane¹³; Florence, Eric¹⁴; Fransen, Katrien¹⁵; Delforge, Marie-Luce¹⁶; Goffard, Jean-Christophe¹⁷; Goubau, Patrick¹⁸ and BREACH, ¹⁹

¹Epidemiology of Infectious Diseases Unit, Scientific Institute of Public Health, Brussels, Belgium. ²AIDS Reference Center, Universitair Ziekenhuis Brussel, Brussels, Belgium. ³AIDS Reference Center, CHU de Liège, Liège, Belgium. ⁴AIDS Reference Laboratory, Universitair Ziekenhuis Brussel, Brussels, Belgium. ⁵AIDS Reference Laboratory, Liège University, Liège, Belgium. ⁶Laboratory of Microbiology, CHU Saint-Pierre, Brussels, Belgium. ⁷AIDS Reference Center, Cliniques Universitaires Saint-Luc, Brussels, Belgium. ⁸AIDS Reference Laboratory, KU Leuven, Leuven, Belgium. ⁹AIDS Reference Center, UZ Leuven, Leuven, Belgium. ¹⁰UZ Gent, AIDS Reference Center, Ghent, Belgium. ¹¹UZ Gent, AIDS Reference Laboratory, Ghent, Belgium. ¹²AIDS Reference Center, CHU de Charleroi, Charleroi, Belgium. ¹³AIDS Reference Center, CHU Saint-Pierre, Brussels, Belgium. ¹⁴AIDS Reference Center, Instituut Tropische Geneeskunde, Antwerp, Belgium. ¹⁵AIDS Reference Laboratory, Instituut Tropische Geneeskunde, Antwerp, Belgium. ¹⁶AIDS Reference Laboratory, University Hospital ULB Erasme, Brussels, Belgium. ¹⁷AIDS Reference Center, University Hospital ULB Erasme, Brussels, Belgium. ¹⁸AIDS Reference Laboratory, Université Catholique de Louvain, Brussels, Belgium. ¹⁹Belgian Research on AIDS and HIV Consortium, Belgium.

Introduction: We studied factors associated with the continuum of HIV care in Belgium.

Methods: Data of the national registration of new HIV diagnosis and of the national cohort of HIV-infected patients in care were combined to obtain estimates of and factors related with proportions of HIV-infected patients in each step of the continuum of care from diagnosis to suppressed viral load (VL). Factors associated with ignorance of HIV seropositivity were analyzed among patients co-infected with HIV and STI in the Belgian STI sentinel surveillance network. Associated factors were identified by multivariate logistic regression.

Results: Among 4038 individuals diagnosed with HIV between 2007 and 2010, 90.3% were linked to care. Of 11684 patients in care in 2010, 90.8% were retained in care up to the following year, 88.3% of those were on ART, of whom 95.3% had suppressed VL (<500 cp/ml) (Figure 1). In multivariate analyses, factors associated with ignoring HIV+ status were being younger ($p < 0.001$), being heterosexual compared to MSM, and of a region of origin other than Belgium, Sub-Saharan Africa and Europe. Non-Belgian regions of origin were associated with lower entry and retention in care ($p < 0.001$ for both). Preoperative HIV testing was associated with lower entry in care ($p = 0.003$). MSM had a higher retention in care ($p < 0.001$), whilst IDU had lower retention ($p = 0.004$). Low CD4 at first clinical contact and clinical reasons for HIV testing were independently associated with being on ART ($p < 0.001$ for both); whilst prenatal HIV diagnosis was associated with lower proportion on ART ($p = 0.016$) and lower proportion with suppressed VL among those on ART ($p = 0.005$). Older age was associated with both being on ART and having suppressed VL among those on ART ($p = 0.007$ and $p < 0.001$ respectively), independently of time since HIV diagnosis (Table 1).

Conclusions: Regions of origin and risk groups (MSM/heterosexual/IDU) are the main factors associated with ignorance of HIV seropositivity, entry and retention in care, but once the HIV patient is retained in care, no effect of these factors on the proportions on ART and with suppressed VL are observed. The association of prenatal HIV diagnosis and proportions on ART and with suppressed VL could be biased by transitory CD4 disturbances during pregnancy and ART discontinuation after pregnancy. The higher probabilities of older patients to be on ART and have suppressed VL once retained in care could be influenced by factors not studied here like comorbidities, adherence or duration on ART.

Published 2 November 2014

Copyright: © 2014 Van Beckhoven D et al; licensee International AIDS Society. This is an Open Access article distributed under the terms of the Creative Commons Attribution 3.0 Unported (CC BY 3.0) License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. The continuum of HIV care in Belgium.

Table 1. Adjusted OR for factors associated with each step of the continuum of HIV care

Risk factors	Adjusted OR (95% CI) ^a Undiagnosed HIV +	Adjusted OR (95% CI) ^a No entry in care	Adjusted OR (95% CI) ^a No retention	Adjusted OR (95% CI) ^a On ART	Adjusted OR (95% CI) ^a Suppressed VL (< 500 cp/ml)
Sex					
Male	1	1	1	1	1
Female	1.32 (0.39–4.44)	0.88 (0.58–1.34)	0.87 (0.70–1.07)	0.76 (0.55–1.04) ^b	1.02 (0.73–1.42)
Age at diagnosis					
< 40 yrs	1	1	1	1	1
≥ 40 yrs	0.42 (0.27–0.64)	0.98 (0.69–1.39)	1.05 (0.86–1.28)	1.31 (1.04–1.65) ^b	1.75 (1.27–2.42)
Way of transmission					
Heterosexual	1	1	1	1	1
MSM	0.39 (0.16–0.95)	0.86 (0.52–1.44)	0.61 (0.47–0.78)	0.80 (0.58–1.09) ^b	1.06 (0.72–1.56)
IDU	/	1.50 (0.55–4.11)	1.88 (1.22–2.88)	2.42 (0.69–8.46) ^b	1.46 (0.52–4.11)
Region of origin					
Belgium	1	1	1	1	1
Sub-Saharan Africa	1.02 (0.33–3.14)	3.11 (1.84–5.26)	1.41 (1.12–1.78)	0.90 (0.66–1.23) ^b	0.73 (0.51–1.03) ^b
Europe	0.83 (0.41–1.69)	2.74 (1.59–4.71)	1.86 (1.38–2.52)	0.97 (0.67–1.40) ^b	0.97 (0.57–1.66) ^b
Other	2.25 (1.26–4.04)	3.23 (1.79–5.83)	1.54 (1.10–2.17)	0.90 (0.59–1.37) ^b	0.89 (0.51–1.48)
Reason for testing					
Patient's request	/	1	1	1	1
Clinical arguments	/	0.98 (0.64–1.50)	0.95 (0.74–1.21)	1.76 (1.34–2.31) ^b	0.92 (0.63–1.35)
Prenatal	/	1.14 (0.51–2.52)	1.16 (0.75–1.78)	0.49 (0.30–0.81) ^b	0.42 (0.23–0.78)
Preoperative	/	3.22 (1.50–6.89)	1.21 (0.72–2.04)	1.14 (0.56–2.31) ^b	0.85 (0.35–2.04)
Other	/	0.92 (0.55–1.54)	1.13 (0.86–1.49)	1.54 (1.11–2.14) ^b	1.03 (0.65–1.63)
CD4 at first visit					
CD4 ≥ 350	/	/	1	1	1
CD4 < 350	/	/	1.16 (0.89–1.51)	8.02 (5.80–11.11)	1.16 (0.78–1.70)

Note: Rem: $p < 0.05$, statistically significant variables presented in italic.

^aAdjusted for sex, age at diagnosis, nationality and ay of transmission; ^badditionally adjusted for CD4 value at first visit.