

Anaesthetic care of patients undergoing primary hip and knee arthroplasty: consensus recommendations from the International Consensus on Anaesthesia-Related Outcomes after Surgery group (ICAROS) based on a systematic review and meta-analysis

Stavros G. Memtsoudis^{1,2,*}, Crispiana Cozowicz^{1,2}, Janis Bekeris^{1,2}, Dace Bekere¹, Jiabin Liu¹, Ellen M. Soffin¹, Edward R. Mariano³, Rebecca L. Johnson⁴, Mary J. Hargett¹, Bradley H. Lee¹, Pamela Wendel¹, Mark Brouillette¹, George Go¹, Sang J. Kim¹, Lila Baaklini¹, Douglas Wetmore¹, Genewoo Hong¹, Rie Goto⁵, Bridget Jivanelli⁵, Eriphyli Argyra⁶, Michael J. Barrington⁷, Alain Borgeat⁸, Jose De Andres^{9,10}, Nabil M. Elkassabany¹¹, Philippe E. Gautier¹², Peter Gerner², Alejandro Gonzalez Della Valle¹, Enrique Goytizolo¹, Paul Kessler¹³, Sandra L. Kopp⁴, Patricia Lavand'Homme¹⁴, Catherine H. MacLean¹⁵, Carlos B. Mantilla⁴, Daniel MacIsaac¹⁶, Alexander McLawhorn¹⁷, Joseph M. Neal¹⁸, Michael Parks¹⁷, Javad Parvizi¹⁹, Lukas Pichler², Jashvant Poeran²⁰, Lazaros A. Poultides²¹, Brian D. Sites²², Otto Stundner², Eric C. Sun²³, Eugene R. Viscusi²⁴, Effrossyni G. Votta-Velis²⁵, Christopher L. Wu¹, Jacques T. Ya Deau¹ and Nigel E. Sharrock¹

¹Department of Anesthesiology, Critical Care and Pain Management, Hospital for Special Surgery and Department of Anesthesiology, Weill Cornell Medical College, New York, NY, USA, ²Department of Anesthesiology, Perioperative Medicine and Intensive Care Medicine, Paracelsus Medical University, Salzburg, Austria, ³Department of Anesthesia, Veterans Affairs Palo Alto Health Care System, Palo Alto, CA, USA, ⁴Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, Minnesota, USA, ⁵Kim Barrett Memorial Library, Hospital for Special Surgery, Weill Cornell Medical College, New York, NY, USA, ⁶Department of Anaesthesiology, Pain and Palliative Care, National and Kapodistrian University of Athens, Athens, Greece, ⁷Department of Medicine & Radiology, The University of Melbourne, Victoria, Australia, ⁸Department of Anesthesiology and Intensive Care Medicine, Universität Zürich, Zurich, Switzerland, ⁹Anesthesia Unit- Surgical Specialties Department, Valencia University Medical School, Spain, ¹⁰Anesthesia, Critical Care, and Pain Management Department, General University Hospital, Valencia, Spain, ¹¹Department of Anesthesiology and Critical Care, University of Pennsylvania, Philadelphia, PA, USA, ¹²Department of Anesthesiology, Clinique Ste-Anne St-Remi, Anderlecht, Belgium, ¹³Department of Anesthesiology, Intensive Care and Pain Medicine, Orthopedic University Hospital, Frankfurt am Main, Germany, ¹⁴Department of Anesthesiology, Cliniques Universitaires Saint-Luc, Brussels, Belgium, ¹⁵Value Management Office, Hospital for Special Surgery, Weill Cornell Medical College, New York, NY, USA, ¹⁶Department of Anesthesiology and Pain Medicine, University of Ottawa, Ottawa, ON, Canada, ¹⁷Department of Orthopedic Surgery, Hip and Knee Replacement, Hospital for Special Surgery, Weill Cornell Medical College, New York, NY, USA, ¹⁸Department of Anesthesiology, Virginia Mason Medical Center, Seattle, WA, USA, ¹⁹Rothman Orthopaedic Institute, Bensalem, PA, USA, ²⁰Department of Population Health Science and Policy, Icahn School of Medicine at Mount Sinai, Institute for Healthcare Delivery Science, New York, NY, USA, ²¹Department of Orthopaedic Surgery, New York Langone Orthopaedic Hospital, New York, NY, USA, ²²Department of Anesthesiology, Dartmouth College Geisel School of Medicine, Hanover, NH, USA, ²³Department of Anesthesia, Stanford University School of Medicine, Stanford, CA, USA, ²⁴Department of Anesthesiology, Pain Center, Sidney Kimmel Medical College at Thomas Jefferson University, Philadelphia, PA, USA and ²⁵Department of Anesthesiology, University of Illinois Hospital and Health Sciences System, Chicago, IL, USA

Abstract

Background: Evidence-based international expert consensus regarding anaesthetic practice in hip/knee arthroplasty surgery is needed for improved healthcare outcomes.

Methods: The International Consensus on Anaesthesia-Related Outcomes after Surgery group (ICAROS) systematic review, including randomised controlled and observational studies comparing neuraxial to general anaesthesia regarding major complications, including mortality, cardiac, pulmonary, gastrointestinal, renal, genitourinary, thromboembolic, neurological, infectious, and bleeding complications. Medline, PubMed, Embase, and Cochrane Library including Cochrane Database of Systematic Reviews, Cochrane Central Register of Controlled Trials, NHS Economic Evaluation Database, from 1946 to May 17, 2018 were queried. Meta-analysis and Grading of Recommendations Assessment, Development and Evaluation approach was utilised to assess evidence quality and to develop recommendations.

Results: The analysis of 94 studies revealed that neuraxial anaesthesia was associated with lower odds or no difference in virtually all reported complications, except for urinary retention. Excerpt of complications for neuraxial vs general anaesthesia in hip/knee arthroplasty, respectively: mortality odds ratio (OR): 0.67, 95% confidence interval (CI): 0.57e0.80/ OR: 0.83, 95% CI: 0.60e1.15; pulmonary OR: 0.65, 95% CI: 0.52e0.80/OR: 0.69, 95% CI: 0.58e0.81; acute renal failure OR: 0.69, 95% CI: 0.59e0.81/OR: 0.73, 95% CI: 0.65e0.82; deep venous thrombosis OR: 0.52, 95% CI: 0.42e0.65/OR: 0.77, 95% CI: 0.64e0.93; infections OR: 0.73, 95% CI: 0.67e0.79/OR: 0.80, 95% CI: 0.76e0.85; and blood transfusion OR: 0.85, 95% CI: 0.82e0.89/OR: 0.84, 95% CI: 0.82e0.87.

Conclusions: Recommendation: primary neuraxial anaesthesia is preferred for knee arthroplasty, given several positive postoperative outcome benefits; evidence level: low, weak recommendation. Recommendation: neuraxial anaesthesia is recommended for hip arthroplasty given associated outcome benefits; evidence level: moderate-low, strong recommendation. Based on current evidence, the consensus group recommends neuraxial over general anaesthesia for hip/ knee arthroplasty.

Trial registry number: PROSPERO CRD42018099935.

Keywords: anaesthesia, epidural; anaesthesia, general; anaesthesia, spinal; arthroplasty, replacement, hip; arthroplasty, replacement, knee; assessment, outcomes

Total joint arthroplasty (TJA) is amongst the most commonly performed surgical procedures in the developed world.¹ Globally, millions of patients receive total hip and knee arthroplasties every year with large projected increases as the population ages.² Despite the fact that TJA represents a value-based solution to end-stage arthritis of the hip and knee,³ the procedure is associated with a moderate risk for

complications. Complications affecting major organ systems have been reported to occur in approximately 8% of patients undergoing either hip or knee arthroplasty.⁴ The identification of risk-modifying perioperative interventions represents an attractive target, given the large burden of resources required for the management of complications on a population-health level.

In this context, a number of recently published population-based studies have supported findings of earlier clinical trials, indicating that the type of anaesthetic technique may influence

Despite this development, definitive evidence in the form of large RCTs or pragmatic, multicentre trials is lacking. Moreover, it is questionable whether such studies will ever exist, given the challenges of feasibility and cost. As population-level data suggesting cost and outcome benefits of neuraxial approaches across a wide range of patient characteristics continue to emerge,^{10e12} it is also unclear if the necessary pre-RCT condition of equipoise can exist to support an experimental trial design.

In light of these factors and given that the utilisation of NA remains low in many countries,¹³ this international group of perioperative clinicians, researchers, quality experts, librarians, educators, and administrators assembled to (i) systematically investigate current published evidence to determine whether the type of anaesthesia technique can influence perioperative outcomes in patients undergoing total hip and knee arthroplasty; (ii) grade the level of evidence quality; and (iii) develop and formulate clinical practice recommendations, each with its own rating of strength.

The aim of the present consensus project was to systematically analyse and interpret current research evidence with regard to the impact of regional, and specifically neuraxial, anaesthesia in comparison to general anaesthesia (GA) on major perioperative outcomes for patients undergoing total hip or knee arthroplasties.

Methods

Consensus group

The International Consensus on Anaesthesia-Related Outcomes after Surgery (ICAROS) consensus group included 50 individuals with extensive expertise in the perioperative care of orthopaedic surgery patients. Included in this multidisciplinary group were anaesthesiologists, orthopaedic surgeons, healthcare outcomes and quality researchers, administrators, librarians, and methodologists from North America, Europe, and Oceania representing 19 nationalities, working in 10 countries. A 10-member steering committee was formed and tasked with overseeing day-to-day aspects of the project.

Study plan and healthcare question

A study plan was specified in advance, defining the healthcare questions and basic parameters, including intervention (NA) and alternative management strategy (GA), population, outcomes of interest, and inclusion and exclusion criteria. The detailed respective protocol, including analyses conducted for this project was registered on the International Prospective Register of Systematic Reviews (protocol number: CRD42018099935).¹⁴ An institutional review board approval was not required because of the analysis of previously published data.

The healthcare questions posed to the group were:

- (i) Does NA vs GA influence perioperative outcomes in patients undergoing total hip arthroplasty (THA)?
- (ii) Does NA vs GA influence perioperative outcomes in patients undergoing total knee arthroplasty (TKA)?

The predefined outcomes of interest included the following major perioperative complications: mortality, cardiac (with and without myocardial infarction), pulmonary (including pneumonia), gastrointestinal, renal (including acute renal failure),

perioperative outcomes.^{5,6} Whilst earlier RCTs suggested a potential benefit of neuraxial anaesthesia (NA) for outcomes, such as blood loss and thromboembolic events, these investigations were not sufficiently powered to study low-incidence outcomes, such as mortality, infectious, or cardiovascular complications.⁷ Furthermore, earlier clinical trials were primarily conducted before the widespread use of chemical thromboembolic prophylaxis and contemporary blood-loss prevention practices.⁸ The advent of population-based scientific approaches utilising large data sets that encompass healthcare information from hundreds of thousands of patients in actual practice environments has allowed researchers to add to the available knowledge in this field. Guided by a series of publications suggesting better outcomes with NA, a number of healthcare entities have developed policies encouraging the use of this anaesthetic type for TJA.⁹

genitourinary (including urinary retention and urinary tract infection), thromboembolic (DVT and pulmonary embolism [PE]), neurological (including CNS complications and stroke), infectious, and wound complications, as well as blood loss (in ml), transfusion requirements (both binary and in ml), and inpatient falls. To account for resource utilisation, the study plan also included outcomes, such as cost of care, length of hospitalisation, and admission to critical care settings. However, because of the lack of studies on cost of care, the outcome could

de facto not be included in the quantitative meta-analysis.^{15,16}

As specified in the study protocol, the consensus group will also address the impact of peripheral nerve block utilisation in patients undergoing total hip and knee arthroplasty. This healthcare question is currently being investigated and will be the focus of a subsequent analysis.

Selection criteria

Eligible studies included RCTs and observational prospective or retrospective studies in adult patients primarily undergoing elective total hip or knee arthroplasties. We included only studies directly comparing perioperative outcomes amongst patients who received NA vs those under GA. GA was defined as total intravenous, inhalational, or combination thereof, or when termed specifically as 'general anaesthesia' by the study authors. NA was defined as spinal, extradural, combined spinal and extradural, and caudal anaesthesia. Exclusion criteria encompassed patients under 18 yr, studies not reporting on postoperative outcomes of interest, case reports, and case series, and also studies without control groups.

Search strategy

A systematic literature search was performed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.

The search strategy, including Medical Subject Headings (MeSH), keywords, and controlled vocabulary terms, was crafted and validated by the expert group in collaboration with two institutional librarians. Medline, PubMed, Embase, and Cochrane Library, including Cochrane Database of Systematic Reviews, Database of Abstracts of Reviews of Effects, Cochrane Central Register of Controlled Trials, Cochrane Methodology Register, Health Technology Assessment Database, and NHS Economic Evaluation Database, were queried from database inception (1946) to May 17, 2018. The search cross-referenced MeSH terms, keywords, and controlled vocabulary terms for the predefined areas of interest according to the healthcare question.

The following is the excerpt of respective search terms: arthroplasty, replacement, hip, total hip arthroplasty, total hip arthroplasties, hip prosthesis, total joint replacement, knee, knee replacement arthroplasty, knee replacement arthroplasties, total knee arthroplasty, knee prosthesis, total knee replacements, lower extremity, lower joints, anaesthesia, neuraxial, spinal, epidural, conduction, regional anaesthesia, intrathecal, peridural, and combined spinal epidural.

The full search strategy is reported in Supplementary materials and can be found in Supplementary Appendix A1. The search yielded 985 studies. In addition to the electronic search, a manual search of previously published corresponding systematic reviews was performed for the purpose of completeness.

Study identification and data extraction

After deduplication, abstracts of 5553 studies were extracted and imported into the Covidence platform. Covidence is a web tool that provides a comprehensive framework for the complete process of a systematic literature review, including the steps of title and abstract screening, full-text review, data extraction, and quality assessment (risk of bias).¹⁷ As required, each step was performed independently by two reviewers. In case of a disagreement, a third reviewer was consulted for resolution.

After the title and abstract screening, full-text articles of 956 studies were imported into Covidence for a detailed review and data extraction. Extracted data were categorised according to the predefined outcomes. Furthermore, within the Covidence platform, the risk of bias for each individual study was assessed and established as high, low, or unknown, according to the Grading of Recommendations Assessment, Development and Evaluation (GRADE) criteria for RCTs and observational studies.¹⁸

A flow chart describing the complete literature search process is depicted in Figure 1.

Quantitative analysis

To provide estimates of intervention effects for each outcome of interest,¹⁹ RCT and observational data were pooled by meta-analysis. Review Manager software (Review Manager (RevMan)

[Computer program]. Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014) was utilised to facilitate data analysis and graphic presentation as is commonly used for preparing Cochrane reviews.²⁰ Summary estimates were calculated separately for each outcome (odds ratios [ORs] and 95% confidence intervals [CIs]), whilst heterogeneity utilising (I^2 statistic) was also determined in quantitative

analysis. For binary outcomes, group-specific risk was presented in events per 1000, whilst the relative effect was presented in ORs. For continuous variables, risk was presented as mean difference. The primary analysis was performed including all eligible studies for both types of surgery, respectively ($n/427$ for TKA; $n/449$ for THA). A separate analysis was performed amongst studies that reported on THA/TKA mixed populations ($n/421$).

Secondary analyses were performed to test the influence of combined NAþGA compared with GA on perioperative outcomes in THA and TKA separately ($n/412$ and $n/44$, respectively), and also in the mixed THA/TKA surgical cohort ($n/48$).

The following are the additional sensitivity analyses:

- Sensitivity analysis to investigate outcomes when only including evidence from RCTs ($n/425$ for THA; $n/412$ for TKA; $n/42$ for THA/TKA)
- Sensitivity analysis to investigate outcomes when removing studies that did not explicitly exclude all revision/trauma-related surgery or bilateral arthroplasties




Fig 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram.

in their cohorts ($n/446$ for THA; $n/425$ for TKA; $n/417$ for THA/TKA)

- (iii) Sensitivity analysis to investigate the potential impact of recent changes in utilisation of perioperative thrombo-embolic prophylaxis protocols on the outcome of thrombo-embolic complications (DVT/PE).

Qualitative analysis

To provide useful recommendations for the practice of evidence-based treatment at the point of care, we utilised the GRADE system.^{15,16} This methodology for rating the quality of evidence and grading the strength of recommendations has been widely adopted for the purpose of providing high-quality summaries of research evidence in systematic reviews and for standardised guideline development. Subsequent to data collection and quantitative analysis, GRADE offers a comprehensive framework for assessing the quality of the body of evidence and for carrying out steps required for developing recommendations.²¹ The concept of the certainty or quality of evidence represents the confidence in effect estimates and the extent to which they are sufficiently credible to support a particular recommendation. GRADE specifies four levels of certainty: high, moderate, low, and very low. This rating is determined for each relevant outcome by the systematic and transparent assessment of study design, limitations of the body of evidence, and special circumstances that increase the quality of evidence. Explicit criteria according to GRADE that were utilised for downgrading the quality of evidence included risk of bias according to study design and study conduct, inconsistency or heterogeneity (lack of similarity of point estimates and overlap of CIs; determination of I^2 statistic), imprecision (optimal information size for adequate power), indirectness (strength of association to the healthcare question), and publication bias (utilising funnel plots).¹⁵ These criteria were assessed for each reported outcome of interest across informing studies. However, risk of bias was also assessed previously for each individual study, whilst in qualitative analysis the impact of risk of bias on cumulative evidence for each outcome was determined. The rationale for upgrading the quality of evidence, particularly for methodologically rigorous observational studies, includes large effect size, presence of a dose-response relationship, or when all plausible confounders or biases would decrease an apparent treatment effect.²² Utilising the GRADEpro software (McMaster University and Evidence Prime Inc.),²³ final results, including the pooled estimates of effect and the quality of evidence, are presented in summary of findings (Tables 1 and 2 for THA and TKA, respectively).

Recommendations

The assessment of the quality of evidence, the formulation of recommendations, and the determination of their strength are separate processes. When moving from evidence to recommendations, the GRADE strategy focuses on integrating factors that are basic for the formulation of guidelines or recommendations.^{19,24} Thus, critical factors beyond the quality of evidence include the balance between benefits and harms; patient values and preferences; resource considerations; and issues pertaining to feasibility, equity, and acceptability of recommendations.¹⁹ GRADE distinguishes between strong and weak recommendations.

The balance between desirable and undesirable outcomes and the application of patients' values determine the direction of the recommendation. Moreover, these factors, along with the quality of evidence, resource implications, and clinical feasibility considerations, determine the strength or grade of recommendations.

Strong recommendations reflect a clear preference for one alternative and should apply to almost all eligible patients. Weak recommendations are appropriate when there is a close balance between desirable and undesirable consequences or alternative management strategies, uncertainty regarding the effects of the alternatives, uncertainty or variability in patient's values and preferences, or questionable cost-effectiveness. Weak recommendations usually require accessing the underlying evidence and a shared decision-making approach.^{15,19,21} In certain circumstances, a strong recommendation is based on low-quality evidence.¹⁶

Modified Delphi process and consensus meeting

Subsequent to analyses completion, two pairs of participants were tasked with summarising the evidence, formulating conclusions, and suggesting recommendations. This work was distributed in the form of white papers for the THA and TKA cohorts separately. The white papers, together with detailed files and summary tables of analysis results, were distributed to the entire group with the request for anonymous edits and comments according to the modified Delphi process,²⁵ and repeated after revisions.²⁶

Finally, the group met in person on December 8, 2018, in New York, NY, USA, to review the process; discuss results; and reach a consensus on conclusions, recommendations, and their strength. Approval was assessed in an anonymous vote after statements were finalised as facilitated by a group discussion.

Results

A summary of findings for patients undergoing THA and TKA, including the estimates of effect and the quality of evidence by outcomes, is found in Tables 1 and 2, respectively.

Additional in-depth quantitative and qualitative analysis data and figures are provided as Supplementary material.

Impact of the type of anaesthesia in total hip arthroplasties

Primary analyses (NA vs GA)

Amongst all hip arthroplasty patients, NA without GA was associated with fewer complications in most categories, except for urinary retention, when compared with patients who received GA (Table 3).

NA was associated with decreased odds for all-cause mortality (OR: 0.67, 95% CI: 0.57, 0.80; absolute effect: 2 per 1000 with GA vs 1 per 1000 with NA, 95% CI: 1, 2), pulmonary complications (OR: 0.65; 95% CI: 0.52, 0.80), pneumonia (OR: 0.69; 95% CI: 0.56, 0.84), and acute renal failure (OR: 0.69; 95% CI: 0.59, 0.81). NA was also associated with fewer thrombo-embolic events compared with GA, including DVT (OR: 0.52; 95% CI: 0.42, 0.65) and PE (OR: 0.63; 95% CI: 0.50, 0.81). Furthermore, CNS complications (OR: 0.39; 95% CI: 0.23, 0.65), stroke (OR: 0.37; 95% CI: 0.22, 0.64), all-cause infections (OR:

Table 1 Grading of Recommendations Assessment, Development and Evaluation (GRADE) summary of findings for total hip arthroplasty (THA). CI, confidence interval; GA, general anaesthesia; MI, myocardial infarction; NA, neuraxial anaesthesia; OR, odds ratio. GRADE Working Group grades of evidence: high certainty (we are very confident that the true effect lies close to that of the estimate of the effect), moderate certainty (we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different), low certainty (our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect), and very low certainty (we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect). *The risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI). ^aPublication bias: funnel plot not symmetric. ^bHeterogeneity: widely differing estimates of effect.

Summary of findings					
NA compared with GA for THA					
Patient or population: THA Setting: perioperative care Intervention: NA Comparison: GA					
Outcomes/complications	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Certainty of the evidence (GRADE)
	Risk with GA	Risk with NA			
Mortality	2 per 1000	1 per 1000 (1e2)	OR: 0.67 (0.57e0.80) (3 RCTs, 4 observational studies)		44OO Low
Cardiac including MI	57 per 1000	53 per 1000 (50e58)	OR: 0.94 (0.88e1.02) (3 RCTs, 3 observational studies)		44OO Low
Cardiac excluding MI	48 per 1000	47 per 1000 (43e50)	OR: 0.96 (0.88e1.03) (2 RCTs, 3 observational studies)		4OOO Very low ^y
MI	3 per 1000	3 per 1000 (2e4)	OR: 0.94 (0.71e1.24) (2 RCTs, 2 observational studies)		44OO Low
Pulmonary	7 per 1000	4 per 1000 (3e5)	OR: 0.65 (0.52e0.80) (3 observational studies)		444O Moderate
Pneumonia	10 per 1000	7 per 1000 (5e8)	OR: 0.69 (0.56e0.84) (2 RCTs, 2 observational studies)		444O Moderate
Gastrointestinal	10 per 1000	8 per 1000 (7e10)	OR: 0.83 (0.67e1.02) (109 732 (1 observational study)		44OO Low
Acute renal failure	15 per 1000	10 per 1000 (9e12)	OR: 0.69 (0.59e0.81) (1 RCT, 5 observational studies)		44OO Low ^z
Urinary retention	111 per 1000	277 per 1000 (199e370)	OR: 3.05 (1.98e4.69) (3 RCTs, 3 observational studies)		444O Moderate
Urinary tract infection	15 per 1000	13 per 1000 (10e15)	OR: 0.86 (0.70e1.06) (2 observational studies)		44OO Low
DVT	15 per 1000	8 per 1000 (6e10)	OR: 0.52 (0.42e0.65) (5 RCTs, 8 observational studies)		444O Moderate
Pulmonary embolism (PE)	3 per 1000	2 per 1000 (2e2)	OR: 0.63 (0.50e0.81) (7 RCTs, 6 observational studies)		44OO Low
Thromboembolism (DVTpPE)	5 per 1000	3 per 1000 (3e4)	OR: 0.61 (0.53e0.71) (15 RCTs, 16 observational studies)		444O Moderate
CNS	2 per 1000	1 per 1000 (0e1)	OR: 0.39 (0.23e0.65) (3 observational studies)		444O Moderate
Stroke	2 per 1000	1 per 1000 (0e1)	OR: 0.37 (0.22e0.64) (2 observational studies)		444O Moderate
	25 per 1000	19 per 1000 (17e20)	OR: 0.73 (0.67e0.79) (2 RCTs, 7 observational studies)		

Continued

Table 1 Continued

Summary of findings					
NA compared with GA for THA					
Patient or population: THA Setting: perioperative care Intervention: NA Comparison: GA					
Outcomes/complications	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Certainty of the evidence (GRADE)
	Risk with GA	Risk with NA			
All infections (including pneumonia and sepsis)					44OO Low
Wound superficial infection	8 per 1000	9 per 1000 (7e12)	OR: 1.21 (0.93e1.56)	(1 RCT, 2 observational studies)	44OO Low
Wound deep infection	7 per 1000	6 per 1000 (5e7)	OR: 0.86 (0.70e1.06)	(3 observational studies)	44OO Low
Blood transfusion	224 per 1000	197 per 1000 (192e205)	OR: 0.85 (0.82e0.89)	(8 RCTs, 9 observational studies)	4OOO Very low ^y
Critical care admission	2 per 1000	1 per 1000 (1e2)	OR: 0.80 (0.49e1.32)	(2 observational studies)	44OO Low
Readmission	38 per 1000	34 per 1000 (30e39)	OR: 0.91 (0.80e1.04)	28 857 (1 observational study)	44OO Low
Nerve injury	2 per 1000	2 per 1000 (1e3)	OR: 0.81 (0.56e1.18)	(1 RCT, 4 observational studies)	44OO Low
Falls	16 per 1000	13 per 1000 (12e15)	OR: 0.81 (0.72e0.92)	166 871 (1 observational study)	44OO Low
Blood loss (ml)	The mean blood loss was 0.	The mean blood loss in the intervention group was 146.12 lower (173.73 lower to 118.51 lower).	d	1546 (12 RCTs, 4 observational studies)	444 O Moderate ^z
Length of stay (days)	The mean length of hospital stay (LOS) was 0.	The mean LOS in the intervention group was 0.16 lower (0.22 lower to 0.1 lower).	d	(1 RCT, 1 observational study)	44OO Low
Blood transfusion (ml)	The mean blood transfusion was 0.	The mean blood transfusion in the intervention group was 187.83 lower (272.29 lower to 103.38 lower).	d	(2 RCTs, 3 observational studies)	44OO Low

Table 2 GRADE summary of findings for total knee arthroplasty (TKA). CI, confidence interval; GA, general anaesthesia; GRADE, Grading of Recommendations Assessment, Development and Evaluation; MI, myocardial infarction; NA, neuraxial anaesthesia; OR, odds ratio. GRADE Working Group grades of evidence: high certainty (we are very confident that the true effect lies close to that of the estimate of the effect), moderate certainty (we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different), low certainty (our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect), and very low certainty (we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect). *The risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI). ¹Imprecision. ²Risk of bias: random sequence generation.

Summary of findings:

NA compared with GA for TKA

Patient or population: TKA
Setting: perioperative care
Intervention: NA Comparison:
GA

Outcomes/complications	Absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with GA	Risk with NA				
Mortality	1 per 1000	1 per 1000 (1e1)	OR: 0.83 (0.60e1.15)	259 847 (2 RCTs, 4 observational studies)	44OO Low	
Cardiac including MI	59 per 1000	60 per 1000 (58e63)	OR: 1.03 (0.98e1.08)	261 695 (1 RCT, 6 observational studies)	44OO Low	
Cardiac excluding MI	57 per 1000	58 per 1000 (55e61)	OR: 1.02 (0.97e1.08)	259 332 (4 observational studies)	44OO Low	
MI	2 per 1000	2 per 1000 (2e3)	OR: 0.99 (0.80e1.22)	261 695 (1 RCT, 6 observational studies)	44OO Low	
Pulmonary	6 per 1000	4 per 1000 (4e5)	OR: 0.69 (0.58e0.81)	259 392 (1 RCT, 4 observational studies)	444O Moderate	
Pneumonia	8 per 1000	6 per 1000 (6e7)	OR: 0.82 (0.72e0.94)	275 947 (1 RCT, 5 observational studies)	44OO Low	
Gastrointestinal	7 per 1000	7 per 1000 (6e8)	OR: 0.99 (0.85e1.15)	223 108 (1 observational study)	44OO Low	
Acute renal failure	14 per 1000	10 per 1000 (9e11)	OR: 0.73 (0.65e0.82)	273 384 (5 observational studies)	44OO Low	
Urinary retention	235 per 1000	203 per 1000 (121e317)	OR: 0.83 (0.45e1.51)	277 (2 RCTs, 1 observational study)	444O Moderate ²	
Urinary tract infection	15 per 1000	12 per 1000 (11e14)	OR: 0.82 (0.71e0.96)	52 779 (4 observational studies)	44OO Low	
DVT	36 per 1000	27 per 1000 (22e32)	OR: 0.77 (0.64e0.93)	19 756 (6 RCTs, 6 observational studies)	44OO Low	
Pulmonary embolism (PE)	6 per 1000	4 per 1000 (4e5)	OR: 0.79 (0.67e0.94)	238 066 (3 RCTs, 4 observational studies)	44OO Low	
Thromboembolism (DVT+PE)	7 per 1000	5 per 1000 (5e6)	OR: 0.77 (0.68e0.88)	257 793 (8 RCTs, 10 observational studies)	44OO Low	
CNS	1 per 1000	1 per 1000 (1e1)	OR: 0.77 (0.55e1.08)	259 594 (1 RCT, 3 observational studies)	44OO Low	
Stroke	1 per 1000	1 per 1000 (1e1)	OR: 0.70 (0.49e1.01)	259 585 (1 RCT, 4 observational studies)	44OO Low	
All infections	22 per 1000	17 per 1000 (16e18)	OR: 0.80 (0.76e0.85)			

Continued

Table 2 Continued

Summary of findings:						
NA compared with GA for TKA						
Patient or population: TKA Setting: perioperative care Intervention: NA Comparison: GA						
Outcomes/complications	Absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with GA	Risk with NA				
				571 503 (1 RCT, 12 observational studies)	44OO Low	
Wound superficial infection	6 per 1000	4 per 1000 (3e6)	OR: 0.77 (0.60e0.98)	52 839 (1 RCT, 4 observational studies)	44OO Low	
Wound deep infection	2 per 1000	2 per 1000 (1e3)	OR: 1.01 (0.60e1.69)	31 843 (3 observational studies)	44OO Low	
Blood transfusion	165 per 1000	142 per 1000 (139e146)	OR: 0.84 (0.82e0.87)	259 332 (4 observational studies)	44OO Low	
Critical care admission	1 per 1000	0 per 1000 (0e1)	OR: 0.17 (0.04e0.75)	20 936 (1 observational study)	444 O Moderate ^y	
Readmission	76 per 1000	38 per 1000 (23e59)	OR: 0.48 (0.29e0.77)	1629 (1 observational study)	444O Moderate	
Nerve injury	4 per 1000	5 per 1000 (2e10)	OR: 1.16 (0.58e2.32)	25 243 (4 observational studies)	44OO Low	
Falls	0 per 1000	0 per 1000 (0e0)	OR: 0.00 (e0.03 to 0.03)	118 (1 observational study)	44OO Low	Not estimable
Blood loss (ml)	The mean blood loss was 0.	The mean blood loss in the intervention group was 13.54 higher (25.75 lower to 52.83 higher).	d	130 (1 RCT)	444O Moderate ^z	
Length of stay (days)	The mean length of hospital stay (LOS) was 0.	The mean LOS in the intervention group was 0.09 lower (0.15 lower to 0.02 lower).	d	36 956 (3 RCTs, 5 observational studies)	44OO Low	

Table 3 Influence of anaesthesia type on perioperative outcomes in total hip arthroplasty. CI, confidence interval; GA, general anaesthesia; MI, myocardial infarction; NA, neuraxial anaesthesia; OR, odds ratio; PE, pulmonary embolism. All infections, including pneumonia and sepsis; pulmonary complications, excluding pneumonia; *n* (NA/GA): total number of patients with NA/GA.

Complication	NA vs GA					NAþGA vs GA				
	Studies	OR (95% CI)	n (NA)	n (GA)	P-value	Studies	OR (95% CI)	n (NA)	n (GA)	P-value
Mortality	Various authors ^{5,27e36}	0.67 (0.57e0.80)	60 499	148 583	<0.0001	Various authors ^{5,30}	0.58 (0.38e0.89)	15 331	98 230	0.014
Cardiac including MI	Various authors ^{5,28,32e34,37}	0.94 (0.88e1.02)	28 182	121 215	0.135	Various authors ^{5,38}	0.76 (0.54e1.07)	15 281	98 139	0.113
Cardiac excluding MI	Various authors ^{5,27,32e34}	0.96 (0.88e1.03)	32 639	133 832	0.255	Memtsoudis and colleagues ⁵	1.01 (0.95e1.09)	15 261	98 122	0.689
MI	Various authors ^{5,33,34,37}	0.94 (0.71e1.24)	23 022	115 759	0.647	Various authors ^{5,38}	0.76 (0.54e1.07)	15 281	98 139	0.113
ulmonary	Various authors ^{5,28,33}	0.65 (0.52e0.80)	28 029	121 058	<0.0001	Memtsoudis and colleagues ⁵	0.66 (0.52e0.84)	15 261	98 122	0.001
Pneumonia	Various authors ^{5,33,34,37}	0.69 (0.56e0.84)	23 022	115 759	<0.0001	Memtsoudis and colleagues ⁵	0.88 (0.74e1.05)	15 261	98 122	0.165
Gastrointestinal	Memtsoudis and colleagues ⁵	0.83 (0.67e1.02)	11 610	98 122	0.078	Memtsoudis and colleagues ⁵	0.79 (0.65e0.95)	15 261	98 122	0.013
Acute renal failure	Various authors ^{5,33,37,39,40}	0.69 (0.59e0.81)	34 366	133 687	<0.0001	Memtsoudis and colleagues ⁵	0.75 (0.65e0.86)	15 261	98 122	<0.0001
Urinary retention	Various authors ^{34,39e43}	3.05 (1.98e4.69)	252	628	<0.0001	Various authors ^{44,45}	1.91 (1.05e3.48)	123	163	0.035
Urinary tract infection	Various authors ^{30,33}	0.86 (0.70e1.06)	11 334	17 648	0.164	Brinker and colleagues ³⁰	1.14 (0.43e2.99)	70	108	0.793
DVT	Various authors ^{30,33,36,41,43,46e53}	0.52 (0.42e0.65)	15 688	20 477	<0.0001	Various authors ^{30,38}	0.81 (0.17e3.89)	90	125	0.795
PE	Various authors ^{5,33e37,41,48e53}	0.63 (0.50e0.81)	34 875	123 934	<0.0001	Memtsoudis and colleagues ⁵	0.68 (0.46e1.03)	15 261	98 122	0.066
DVTþPE	Various authors ^{5,30,33e37,41,43,46e57}	0.61 (0.53e0.71)	59 573	157 731	<0.0001	Various authors ^{5,30,38}	0.69 (0.47e1.02)	15 351	98 247	0.065
CNS	Various authors ^{5,33,58}	0.39 (0.23e0.65)	22 977	115 712	<0.0001	Various authors ^{5,34,59}	0.68 (0.42e1.09)	15 306	98 162	0.112
Stroke	Various authors ^{5,33}	0.37 (0.22e0.64)	22 927	115 662	<0.0001	Memtsoudis and colleagues ⁵	0.71 (0.44e1.16)	15 261	98 122	0.176
All infections	Various authors ^{5,28,30,33,34,37}	0.73 (0.67e0.79)	62 385	254 465	<0.0001	Various authors ^{5,30}	0.86 (0.79e0.92)	30 592	196 352	<0.0001
Wound (superficial)	Various authors ^{30,33,34}	1.21 (0.93e1.56)	11 363	17 679	0.152	Brinker and colleagues ³⁰	1.56 (0.21e11.33)	70	108	0.661
Wound (deep)	Various authors ^{28,33,54}	0.86 (0.70e1.06)	24 603	35 688	0.159					
Blood transfusion	Various authors ^{5,30,33,34,37,41,43,60e69}	0.85 (0.82e0.89)	25 033	117 443	<0.0001	Various authors ^{5,30,38,61}	0.78 (0.75e0.82)	15 421	98 317	<0.0001
Critical care	Various authors ^{27,33}	0.80 (0.49e1.32)	20 690	33 125	0.387					
Readmission	Haughom and colleagues ³³	0.91 (0.80e1.04)	11 317	17 540	0.161					
Nerve injury	Various authors ^{30,33,34,69}	0.81 (0.56e1.18)	19 842	27 106	0.278	Brinker and colleagues ³⁰	0.30 (0.01e6.39)	70	108	0.442
Falls	Kendrick and colleagues ⁷¹	0.81 (0.72e0.92)	20 985	145 886	0.001					
Blood loss (ml)	Various authors ^{36,43,50e52,60e62,66,68,72e77}	e146.12 (e173.73 to e118.51)	902	644	<0.0001	Various authors ^{38,59,61,74,77e79}	e20.13 (e50.10 to 9.83)	226	216	0.188
Length of stay (days)	Various authors ^{28,80}	e0.16 (e0.22 to e0.10)	5146	5442	<0.0001	Benson and colleagues ⁵⁹	e6.00 (e14.77 to 2.77)	16	9	0.18
Blood transfusion (ml)	Various authors ^{43,50,51,60,66}	e187.83 (e272.29 to e103.38)	310	195	<0.0001					

0.73; 95% CI: 0.67, 0.79), blood transfusion requirements (OR: 0.85; 95% CI: 0.82, 0.89), and postoperative falls (OR: 0.81; 95% CI: 0.72, 0.92) were reduced with NA vs GA.

We did not identify any differences in cardiac, gastroin- testinal, or wound complications; critical care admissions; readmissions; and nerve injuries between NA and GA amongst hip arthroplasty patients.

Impact of the type of anaesthesia in total knee arthroplasties

Primary analyses (NA vs GA)

Amongst patients undergoing total knee arthroplasties, the utilisation of NA in comparison to GA was associated with improved outcomes with regard to multiple complications (Table 4). Amongst patients who received NA, reduced odds were observed for pulmonary complications (OR: 0.69; 95% CI: 0.58e0.81), pneumonia (OR: 0.82; 95% CI: 0.72, 0.94), acute renal failure (OR: 0.73; 95% CI: 0.65, 0.82), urinary tract infection (OR: 0.82; 95% CI: 0.71, 0.96), DVT (OR: 0.77; 95% CI: 0.64, 0.93), PE (OR: 0.79; 95% CI: 0.67, 0.94), all-cause infections (OR: 0.80; 95% CI: 0.76, 0.85), superficial wound infections (OR: 0.77; 95% CI: 0.60, 0.98), blood transfusions (OR: 0.84; 95% CI: 0.82, 0.87), critical care admissions (OR: 0.17; 95% CI: 0.04, 0.75), and readmissions (OR: 0.48; 95% CI: 0.29, 0.77).

Impact of the type of anaesthesia in studies reporting outcomes in mixed total knee/hip arthroplasties

Primary analyses (NA vs GA)

The results are presented in Supplementary Table A5. Overall, improved outcomes were seen in association with the use of NA vs GA in this cohort of studies.

Secondary analyses (NAþGA vs GA)

In a secondary analysis, we compared the utilisation of combined NAþGA vs GA only to assess the impact on studied outcomes in patients undergoing THA and TKA (Tables 3 and 4, and Supplementary Table A5). The output indicated a similar trend as observed in the NA vs GA analysis. The out- comes with significantly reduced odds for combined NAþGA vs GA included mortality, pulmonary complications, gastro- intestinal complications, acute renal failure, all-cause in- fections, and blood transfusions, whilst the odds for urinary retention were increased as seen in the NA vs GA comparison.

Sensitivity analyses

Randomised clinical trials only

The first sensitivity analysis focused on RCTs only and verified that NA was associated with fewer thromboembolic events than GA (Tables 5 and 6, and Supplementary Table A6). NA patients also had less blood loss and received lower blood transfusion volumes (Table 5). This analysis did not present statistically significant differences in other complications, which may be attributable to the much smaller sample size in RCTs compared with population-based analyses.

Exclusion of studies likely containing a minority of revision/trauma surgery or bilateral arthroplasty cases

Our primary analysis included all patients from all candidate studies, which encompassed RCTs and observational studies. In some of these investigations, revision/trauma-related arthroplasty patients could not be excluded with certainty. To test the potential effect that this patient population may have on outcomes, we excluded them in a sensitivity analysis. The relationship between anaesthetic type and outcomes when excluding revision/trauma arthroplasty was nearly identical compared with the primary inclusive analysis (Supplementary Tables A2eA4).

Sensitivity analysis: thromboembolic complications (DVTþPE)

To account for potential prognostic imbalance as a result of recent emerging differences in perioperative care with regard to the implementation of thrombosis prophylaxis in recent years, we performed a further sensitivity analysis. Estimates of inter- vention effects were established for the outcome of thrombo- embolic complications (DVTþPE) when including all eligible studies, when excluding studies without thrombosis prophy- laxis, and when excluding studies published before 1995.

NA was associated with a 24% reduction in thromboem- bolic events when including all studies (n /437; OR: 0.76; 95% CI: 0.71, 0.83), a 14% reduction when excluding studies lacking thromboembolic prophylaxis (n /49; OR: 0.86; 95% CI: 0.79, 0.92), and a 16% reduction when excluding studies before 1995 (n /414; OR: 0.84; 95% CI: 0.78, 0.90).

Discussion

Recommendations and comments

Does type of anaesthesia influence perioperative outcomes in THA?

The utilisation of NA over GA for THA was associated with lower complication odds for most studied outcomes. The uti- lisation of combined NA and GA was also associated with better perioperative outcomes compared with GA alone, although the magnitude and diversity of benefits were decreased compared with using NA alone (Tables 1 and 3).

- (i) Level of evidence: low to moderate
- (ii) Recommendation: NA is recommended for primary uni- lateral THA when there is no significant contraindication or special circumstance to preclude its use.
- (iii) Strength of recommendation: strong
- (iv) Rationale: Based on the findings of our analysis and the grading of evidence, the group reached a unanimous de- cision on the aforementioned recommendation. The re- sults of all analyses showed improvement in outcomes with NA compared with GA in most cases, or no impact, with the sole exception of urinary retention, albeit the latter is a known, expected side-effect of NA.¹⁰¹

The level of evidence underlying the individual analyses by outcome was low to moderate. When considering the factors integrated by the GRADE approach for the development of recommendations,¹⁹ the majority of the group (n /433 out of 43 votes) determined it to be overall strong.

The latter conclusion was based on the observations that:

- (i) the evidence was largely in favour of the intervention, (ii)

Table 4 Influence of anaesthesia type on perioperative outcomes in total knee arthroplasty. CI, confidence interval; GA, general anaesthesia; MI, myocardial infarction; NA, neuraxial anaesthesia; OR, odds ratio; PE, pulmonary embolism. All infections, including pneumonia and sepsis; pulmonary complications, excluding pneumonia; *n* (NA/GA): total number of patients with NA/GA.

Complication	NA vs GA					NA vs GA				
	Studies	OR (95% CI)	n (NA)	n (GA)	P-value	Studies	OR (95% CI)	n (NA)	n (GA)	P-value
Mortality	Various authors ^{5,6,54,57,81,82}	0.83 (0.60e1.15)	43 653	216 194	0.261	Memtsoudis and colleagues ⁵	0.73 (0.51e1.05)	34 135	194 682	0.094
Cardiac including MI	Various authors ^{5,6,54,57,81,83,84}	1.03 (0.98e1.08)	44 831	216 864	0.324	Various authors ^{5,85}	1.07 (0.85e1.34)	34 165	194 715	0.553
Cardiac excluding MI	Various authors ^{5,6,54,57}	1.02 (0.97e1.08)	43 386	215 946	0.349	Memtsoudis and colleagues ⁵	1.07 (1.02e1.12)	34 135	194 682	0.007
MI	Various authors ^{5,6,54,57,81,83,84}	0.99 (0.80e1.22)	44 831	216 864	0.896	Various authors ^{5,85}	1.07 (0.85e1.34)	34 165	194 715	0.553
Pulmonary	Various authors ^{5,6,54,57,86}	0.69 (0.58e0.81)	43 416	215 976	<0.0001	Various authors ^{5,85}	0.89 (0.77e1.03)	34 165	194 715	0.132
Pneumonia	Various authors ^{5,6,54,57,86,87}	0.82 (0.72e0.94)	50 804	225 143	0.003	Memtsoudis and colleagues ⁵	1.02 (0.90e1.16)	34 135	194 682	0.727
Gastrointestinal	Memtsoudis and colleagues ⁵	0.99 (0.85e1.15)	28 426	194 682	0.855	Various authors ^{5,85}	1.07 (0.93e1.22)	34 165	194 715	0.344
Acute renal failure	Various authors ^{5,6,54,57}	0.73 (0.65e0.82)	49 416	223 968	<0.0001	Memtsoudis and colleagues ⁵	0.96 (0.87e1.05)	34 135	194 682	0.377
Urinary retention	Various authors ^{32,86,88}	0.83 (0.45e1.51)	111	166	0.537					
Urinary tract infection	Various authors ^{6,54,57,87}	0.82 (0.71e0.96)	22 348	30 431	0.011					
DVT	Various authors ^{83,89,90,92e95}	0.77 (0.64e0.93)	9466	10 222	0.005	⁸⁵	0.53 (0.05e6.21)	30	33	0.617
PE	Various authors ^{5,6,42,82,90,92,94}	0.79 (0.67e0.94)	34 890	203 176	0.007	Memtsoudis and colleagues ⁵	0.78 (0.66e0.93)	34 135	194 682	0.006
DVT vs PE	Various authors ^{5,6,42,47,56,82,}	0.77 (0.68e0.88)	44 373	213 420	<0.0001	Various authors ^{5,85}	0.78 (0.66e0.93)	34 165	194 715	0.005
CNS	Various authors ^{5,6,54,57,81}	0.77 (0.55e1.08)	43 520	216 074	0.133	Various authors ^{5,85,96}	1.03 (0.75e1.43)	34 270	194 823	0.84
Stroke	Various authors ^{5,6,54,57,82}	0.70 (0.49e1.01)	43 519	216 066	0.059	Various authors ^{5,85}	1.06 (0.76e1.49)	34 165	194 715	0.72
All infections	Various authors ^{5,6,54,57,86,87}	0.80 (0.76e0.85)	109 150	462 353	<0.0001	Memtsoudis and colleagues ⁵	0.98 (0.93e1.03)	68 270	389 364	0.464
Wound (superficial)	Various authors ^{6,54,57,86,87}	0.77 (0.60e0.98)	22 378	30 461	0.034					
Wound (deep)	Various authors ^{6,57,87}	1.01 (0.60e1.69)	14 164	17 679	0.982					
Blood transfusion	Various authors ^{5,6,54,57}	0.84 (0.82e0.87)	43 386	215 946	<0.0001	Memtsoudis and colleagues ⁵	1.02 (0.99e1.05)	34 135	194 682	0.197
Critical care	Basques and colleagues ⁵⁴	0.17 (0.04e0.75)	8184	12 752	0.019					
Readmission	Belmont and colleagues ⁹⁷	0.48 (0.29e0.77)	586	1043	0.003					
Nerve injury	Various authors ^{6,57,70}	1.16 (0.58e2.32)	7304	17 939	0.665					
Falls	Harsten and colleagues ³²	0.00 (0.00e0.00)	58	60	<0.0001	Kendrick and colleagues ⁷¹	0.91 (0.81e1.02)	24 699	145 886	0.092
Blood loss (ml)	Zhou and colleagues ⁹⁵	13.54 (e25.75 to 52.83)	63	67	0.499	Kudoh and colleagues ⁹⁶	13.10 (e18.99 to 45.19)	75	75	0.424
Length of stay (days)	Various authors ^{6,54,57,81,82,98e100}	e0.09 (e0.15 to e0.02)	15 326	21 630	0.009					

Table 5 Subgroup RCTs: influence of anaesthesia type on perioperative outcomes in total hip arthroplasty. CI, confidence interval; GA, general anaesthesia; MI, myocardial infarction; NA, neuraxial anaesthesia; OR, odds ratio; PE, pulmonary embolism. All infections, including pneumonia and sepsis; pulmonary complications, excluding pneumonia; *n* (NA/GA): total number of patients with NA/GA.

Complication	NA vs GA					NApGA vs GA				
	Studies	OR (95% CI)	n (NA)	n (GA)	P-value	Studies	OR (95% CI)	n (NA)	n (GA)	P-value
Mortality	Various authors ^{32,34,36}	0.34 (0.01e8.80)	135	137	0.519					
Cardiac including MI	Various authors ^{32,34,37}	0.82 (0.19e3.48)	153	157	0.783	Dauphin and colleagues ³⁸	1.78 (0.15e21.51)	20	17	0.651
Cardiac excluding MI	Various authors ^{32,34}	0.65 (0.08e5.38)	87	91	0.687	Dauphin and colleagues ³⁸	1.78 (0.15e21.51)	20	17	0.651
	Various authors ^{34,37}	0.75 (0.14e4.07)	95	97	0.736					
Pneumonia	Various authors ^{34,37}	1.03 (0.14e7.53)	95	97	0.973					
Acute renal failure	Liang and colleagues ³⁷	0.33 (0.01e8.21)	66	66	0.498					
Urinary retention	Various authors ^{34,41,42}	1.65 (0.89e3.05)	158	162	0.113					
DVT	Various authors ^{36,41,50,51}	0.33 (0.20e0.55)	177	174	<0.0001	Dauphin and colleagues ³⁸	0.81 (0.17e3.89)	20	17	0.795
PE	Various authors ^{34,36,37,41,50,51}	0.40 (0.20e0.79)	255	257	0.008					
DVTpPE	Various authors ^{34,36,37,41,50,51,55,56}	0.43 (0.30e0.63)	482	479	<0.0001	Dauphin and colleagues ³⁸	0.81 (0.17e3.89)	20	17	0.795
CNS						Various authors ^{34,59}	0.26 (0.03e2.28)	45	40	0.222
All infections	Various authors ^{34,37}	1.03 (0.14e7.53)	95	97	0.973					
Wound (superficial)	³⁴	0.33 (0.03e3.40)	29	31	0.354					
Blood transfusion	Various authors ^{34,37,41,61,63,67,68}	0.43 (0.28e0.65)	357	364	<0.0001	Various authors ^{38,61}	0.50 (0.24e1.05)	90	87	0.067
Nerve injury	Hole and colleagues ³⁴	0.34 (0.01e8.80)	29	31	0.519					
Blood loss (ml)	Various authors ^{36,50,51,61,62,68,72e75,77}	e121.82 (e152.22 to e91.42)	334	335	<0.0001	Various authors ^{38,59,61,74,77e79}	e20.13 (e50.10 to 9.83)	226	216	0.188
Length of stay (days)	Williams-Russo and colleagues ⁸⁰	e3.00 (e6.25 to 0.25)	44	46	0.07	Benson and colleagues ⁵⁹	e6.00 (e14.77 to 2.77)	16	9	0.18
Blood transfusion (ml)	Various authors ^{50,51}	e542.64 (e771.95 to e313.32)	45	45	<0.0001					

Table 6 Subgroup RCTs: influence of anaesthesia type on perioperative outcomes in total knee arthroplasty. CI, confidence interval; GA, general anaesthesia; MI, myocardial infarction; NA, neuraxial anaesthesia; OR, odds ratio; PE, pulmonary embolism. All infections, including pneumonia and sepsis; pulmonary complications, excluding pneumonia; *n* (NA/GA): total number of patients with NA/GA.

Complication	NA vs GA					NA+GA vs GA				
	Studies	OR (95% CI)	n (NA)	n (GA)	P-value	Studies	OR (95% CI)	n (NA)	n (GA)	P-value
Mortality	Various authors ^{81,82}	0.93 (0.13e6.64)	267	248	0.941					
Cardiac including MI	Williams-Russo and colleagues ⁸¹	1.28 (0.28e5.84)	134	128	0.748					
MI	Williams-Russo and colleagues ⁸¹	0.95 (0.19e4.82)	134	128	0.955					
Pulmonary	Chu and colleagues ⁸⁶	0.48 (0.04e5.63)	30	30	0.561					
Pneumonia	Chu and colleagues ⁸⁶	0.19 (0.01e4.06)	30	30	0.286					
Urinary retention	Various authors ^{32,86}	0.86 (0.47e1.59)	88	90	0.628					
DVT	Various authors ^{56,82,89e91,95}	0.82 (0.56e1.18)	256	327	0.283					
PE	Various authors ^{42,82,90}	1.17 (0.45e3.03)	163	149	0.748					
DVT+PE	Various authors ^{42,56,82,89,90,95}	0.78 (0.56e1.10)	436	498	0.157					
CNS	Williams-Russo and colleagues ⁸¹	1.31 (0.59e2.89)	134	128	0.503	Kudoh and colleagues ⁹⁶	0.74 (0.16e3.42)	75	75	0.7
Stroke	Williams-Russo and colleagues ⁸²	2.73 (0.11e67.61)	133	120	0.54					
All infections	Chu and colleagues ⁸⁶	0.19 (0.01e4.06)	30	30	0.286					
Wound (superficial)	Chu and colleagues ⁸⁶	0.48 (0.04e5.63)	30	30	0.561					
Falls	Harsten and colleagues ³²	0.00 (0.00e0.00)	58	60	<0.0001					
Blood loss (ml)	Zhou and colleagues ⁹⁵	13.54 (e25.75 to 52.83)	63	67	0.499	Kudoh and colleagues ⁹⁶	13.10 (e18.99 to 45.19)	75	75	0.424
Length of stay (days)	Various authors ^{81,82,98}	e0.14 (e0.56 to 0.28)	308	295	0.512					

the desirable effects of the intervention outweigh the undesirable ones, (iii) the intervention was associated with neutral to beneficial resource utilisation, (iv) the intervention is acceptable to stakeholders, and (v) the intervention is feasible.

Does type of anaesthesia influence perioperative outcomes in TKA?

Compared with GA, NA was associated with fewer complications or no difference in complications in all reported outcomes after TKA (Tables 2 and 4).

NA was associated with lower odds of thromboembolic events and blood transfusion, and also infectious complications, including pneumonia and all-cause infections. Furthermore, lower odds for acute renal failure and respiratory complications were found amongst patients receiving NA for TKA. With regard to outcomes of resource utilisation, NA was associated with fewer admissions to critical care units, lower rates of hospital readmissions, and a shorter length of hospital stay (mean difference: e0.08; 95% CI: e0.15 to 0.01 days).

Our analysis failed to find any significant differences in the odds for mortality, composite CNS complications, or stroke. There was also no effect of anaesthetic type on cardiac or gastrointestinal complications.

- (i) Level of evidence: low
- (ii) Recommendation: Provided no contraindication, a primary neuraxial anaesthetic technique is preferred for TKA, given several positive benefits of NA on important post-TKA outcomes, together with no evidence of worse outcomes.
- (iii) Strength of recommendation: weak
- (iv) Rationale: Based on the findings of our analysis and the grading of the level of evidence, the group reached a majority ($n/42$ out of 43 votes) decision on the aforementioned recommendation. The results of all analyses showed improvement with NA for outcomes compared with GA for some but not all outcomes. The effect was smaller than that seen in the larger THA cohort.

The level of evidence underlying the individual analyses by outcome was low.

When considering the factors integrated by the GRADE approach for the development of recommendations, the majority of the group ($n/43$ out of 43 votes) determined it to be overall weak.

The latter conclusion was based on the observations that the evidence was in favour of the intervention, but to a lesser extent than that observed in the THA cohort. However, the group believed that the desirable effects of the intervention outweigh the undesirable effects, and that the intervention was associated with beneficial resource utilisation. Further, the intervention is acceptable to stakeholders and is clinically feasible.

Comments

Several limitations to our consensus approach have to be considered. Perioperative care has evolved significantly over years and decades, including surgical techniques. This may be a source of unmeasured or unknown confounding that is not adequately balanced by randomisation.

Further, the group discussed extensively the lack of detailed information regarding the potentially wide variability in the conduct of GA and the potential influence of GA technique on outcomes. Whilst NA as a technique may vary to

certain degrees (type of local anaesthetic used, use of spinal vs extradural vs combined spinal/extradural, and level of neuraxial block), the group agreed that the conduct of the technique and its major characteristics are standardised and have been in place for many decades. In contrast, the conduct of GA has evolved significantly over time with changes in pharmacological agents (both intravenous and inhalational), assistive technology (target-controlled infusion), airway devices, monitoring, and ventilation equipment, and also care strategies.

Therefore, it seems appropriate to re-evaluate the differential impact of modern general anaesthetic techniques in this context once such granular information becomes reliably available in the future.

Additional factors that may influence outcomes include the use of procedural sedation and its depth, which may, in practice, approach levels seen with GA.¹⁰² However, at this time, such an analysis is not feasible because of the lack of adequate data. In addition, the inherent anaesthetic-related risks of each technique (GA or NA) were not considered in this analysis, but are rare for either approach.

In the last decade, advances in regional anaesthesia, such as the utilisation of ultrasound-guided peripheral nerve block techniques, have gained significant popularity in the clinical setting. Thus, our research group is currently reviewing evidence regarding the perioperative impact of peripheral nerve blocks. However, given the numerous options and combinations of various anaesthesia-related procedures, further studies are needed to address specifically the impact of peripheral nerve blocks as adjuncts to GA when compared with NA.

Further, the group discussed what future research would be needed to derive definitive data on the questions addressed in this consensus article. Whilst large, multicentre RCTs or pragmatic trials may provide definitive evidence, they are not and may never be available. Future studies are indicated to better evaluate the mechanisms by which the observed beneficial effects associated with NA are realised. The group acknowledged that, whilst a plausible mechanism for improved outcomes is likely related to NA-associated reductions in stress response,

the body of evidence establishing this link is scarce.¹⁰³ Further, it was determined that future research is needed to elucidate the relationship between anaesthetic type and outcomes in the ever-increasing commonality of high-risk patient populations presenting for joint arthroplasty. Moreover, comparative literature for some complications, such as postoperative cognitive dysfunction, is rare, and these topics require more scientific investigations to allow robust analysis and conclusions in the

context of anaesthesia practice.^{58,81} Finally, the group commented that, given the potential benefits and relative under-utilisation of NA, research with focus on identification and amelioration of barriers to the widespread implementation of NA techniques is needed.

Executive summary

Does type of anaesthesia influence perioperative outcomes in THA?

The utilisation of NA over GA for THA was associated with lower complication risk for most studied outcomes. Furthermore, the utilisation of combined NA and GA was also associated with better perioperative outcomes compared with GA alone, although the magnitude and diversity of benefits were decreased compared with using NA alone (Tables 1 and 3).

- (i) Level of evidence: low to moderate
- (ii) Recommendation: NA is recommended for primary uni- lateral THA when there is no significant contraindication or special circumstance to preclude its use.
- (iii) Strength of recommendation: strong

Does type of anaesthesia influence perioperative outcomes in TKA?

- (i) Level of evidence: low
- (ii) Recommendation: Provided no contraindication, a primary neuraxial anaesthetic technique is preferred for TKA, given several positive benefits of NA on important post-TKA outcomes, together with no evidence of worse outcomes.
- (iii) Strength of recommendation: weak

Authors' contributions

Study conception: SGM

Study design/planning/execution: SGM, NES, CC, JB, JL, EMS, ERM, RLJ, MJH, GG

Reviewing/expanding study plan: EA, MJB, AB, JDA, NE, PEG, PG, AGDV, EG, PK, SLK, PL'H, CHML, CBM, DM, AM, JMN, MP, JPa, LP, JPo, LAP, BDS, OS, ECS, ERV, EGV-V, CW, JTYD Literature search: CC, RG, BJ, LP, BHL, PW, MB, GG, SJK, LB, DW, GH Data extraction: JB, DB, CC, BHL, PW, MB, GG, SJK, LB, DW, GH, JL, SGM

Data analysis: CC, SGM, NES, JPo, JB, JL, ES, ERM, RLJ, MJH, GG Reviewing results of data analysis: EA, MJB, AB, JDA, NE, PEG, PG, AGDV, EG, PK, SLK, PL'H, CHML, CBM, DM, AM, JMN, MP, JPa, LP, JPo, LAP, BDS, OS, ECS, ERV, EGV-V, CLW, JTYD Interpreting results: CC, SGM, NES, JPo, JB, JL, ES, ERM, RLJ, MJH, GG

Reviewing/editing white papers: EA, MJB, AB, JDA, NE, PEG, PG, AGDV, EG, PK, SLK, PL'H, CHML, CBM, DM, AM, JMN, MP, JPa, LP, JPo, LAP, BDS, OS, ECS, ERV, EGV-V, CLW, JTYD Writing paper: SGM, CC, NES, JB, JL, EMS, ERM, RLJ, MJH, GG

All authors reviewed, commented on, and approved the study plan; reviewed the data and the analysis results; commented on and gave feedback to the interpretation of results, including quantitative and qualitative analyses; and convened in person or were given the opportunity to join remotely in an all-day consensus conference held on December 8, 2018 at the Hospital for Special Surgery, New York, NY, USA, where the entire analysis steps and results were presented, and the GRADE approach was utilised for the interpretation of the body of evidence and the formation of recommendations.

SGM had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Declarations of interest

SGM is a director of the boards of the American Society of Regional Anesthesia and Pain Medicine and the president of the Society of Anesthesia and Sleep Medicine. He is a one-time consultant for Teikoku, Sandoz Inc. and a consultant/investor for HATH. Furthermore, SGM has a US Patent application pending for a Multicatheter Infusion System (US-2017- 0361063). He is the owner of SGM Consulting, LLC, and co- owner of FC Monmouth, LLC. None of these relations influenced the conduct of the present project. ERM is a director of

the board of the American Society of Regional Anesthesia and Pain Medicine and an officer of the California Society of Anesthesiologists. ERM is also an employee of the United States government, and his contribution to this project is supported with resources based at the Veterans Affairs (VA) Palo Alto Health Care System (Palo Alto, CA, USA). The contents do not represent the views of VA or the United States Government. NE is a board member of the American Society of Regional Anesthesia and Pain Medicine. NE is also a consultant for Foundry Therapeutics, but declared no conflict of interest. ECS reports consulting fees from Egalet, Inc. and the Mission Lisa foundation and acknowledges funding from the National Institute on Drug Abuse (K08DA042314), which are unrelated to this work. The other authors declare that they have no conflicts of interest.

Funding

Department of Anesthesiology, Critical Care, and Pain Management, Hospital for Special Surgery, New York, NY, USA.

Disclaimer

The conclusions and recommendations resulting from this project are not intended to establish practice guidelines or standards, nor can they if followed guarantee successful outcomes. Many adequate reasons exist why a clinician or patient may deviate from the recommendations in this article, including, but not limited to, medical circumstances, individual patient and clinician preferences, and the availability of resources. The present conclusions and recommendations are based on the currently available literature, established in a systematic review process; thus, reassessment and revisions are required as new or different evidence emerges.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bja.2019.05.042>.

References

1. Pabinger C, Geissler A. Utilization rates of hip arthroplasty in OECD countries. *Osteoarthritis Cartilage* 2014; 22: 734e41
2. Kurtz S, Ong K, Lau E, Mowat F, Halpern M. Projections of primary and revision hip and knee arthroplasty in the United States from 2005 to 2030. *J Bone Jt Surg Am* 2007; 89: 780e5
3. Ethgen O, Bruyere O, Richy F, Dardennes C, Reginster JY. Health-related quality of life in total hip and total knee arthroplasty. A qualitative and systematic review of the literature. *J Bone Jt Surg Am* 2004; 86: 963e74
4. Memtsoudis SG, Ma Y, Gonzalez Della Valle A, et al. Demographics, outcomes, and risk factors for adverse events associated with primary and revision total hip arthroplasties in the United States. *Am J Orthop* 2010; 39: E72e7
5. Memtsoudis SG, Sun X, Chiu YL, et al. Perioperative comparative effectiveness of anesthetic technique in orthopedic patients. *Anesthesiology* 2013; 118: 1046e58
6. Pugely AJ, Martin CT, Gao Y, Mendoza-Lattes S, Callaghan JJ. Differences in short-term complications

- between spinal and general anesthesia for primary total knee arthroplasty. *J Bone Jt Surg Am* 2013; 95: 193e9
7. Mauermann WJ, Shilling AM, Zuo Z. A comparison of neuraxial block versus general anesthesia for elective total hip replacement: a meta-analysis. *Anesth Analg* 2006; 103: 1018e25
 8. Johnson R, Kopp S, Burkle C, et al. Neuraxial vs general anaesthesia for total hip and total knee arthroplasty: a systematic review of comparative-effectiveness research. *Br J Anaesth* 2016; 116: 163e76
 9. Health Quality Ontario & Ministry of Health and Long- Term Care. *Quality-based procedures: clinical handbook for primary hip and knee replacement* 2014. 1e95, http://www.health.gov.on.ca/en/pro/programs/ecfa/docs/qbp_prihipknee.pdf. [Accessed 7 July 2019]
 10. Memtsoudis SG, Poeran J, Zubizarreta N, et al. Do hospitals performing frequent neuraxial anesthesia for hip and knee replacements have better outcomes? *Anesthesiology* 2018; 129: 428e39
 11. Memtsoudis SG, Poeran J, Zubizarreta N, Rasul R, Opperer M, Mazumdar M. Anesthetic care for orthopedic patients: is there a potential for differences in care? *Anesthesiology* 2016; 124: 608e23
 12. Memtsoudis SG, Rasul R, Suzuki S, et al. Does the impact of the type of anesthesia on outcomes differ by patient age and comorbidity burden? *Reg Anesth Pain Med* 2014; 39: 112e9
 13. Cozowicz C, Poeran J, Memtsoudis SG. Epidemiology, trends, and disparities in regional anaesthesia for orthopaedic surgery. *Br J Anaesth* 2015; 115: ii57e67
 14. *International Prospective Register of Systematic Reviews*. <https://www.crd.york.ac.uk/prospero/>. [Accessed 7 July 2019]
 15. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008; 336: 924e6
 16. Balshem H, Helfand M, Schünemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011; 64: 401e6
 17. Covidence systematic review software, Veritas Health Innovation, Melbourne, Australia. Available at www.covidence.org (please see <https://support.covidence.org/help/how-can-i-cite-covidence>)
 18. Guyatt GH, Oxman AD, Vist G, et al. GRADE guidelines: 4. Rating the quality of evidenced study limitations (risk of bias). *J Clin Epidemiol* 2011; 64: 407e15
 19. Neumann I, Santesso N, Akl EA, et al. A guide for health professionals to interpret and use recommendations in guidelines developed with the GRADE approach. *J Clin Epidemiol* 2016; 72: 45e55
 20. *Cochrane Community. RevMan 5*. <https://community.cochrane.org/help/tools-and-software/revman-5>. [Accessed 7 July 2019]
 21. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction and GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011; 64: 383e94
 22. Guyatt GH, Oxman AD, Sultan S, et al. GRADE guidelines: 9. Rating up the quality of evidence. *J Clin Epidemiol* 2011; 64: 1311e6
 23. McMaster University. *GRADEpro GDT: GRADEpro Guideline Development Tool [software]* 2015. <https://grade.pro.org/>. [Accessed 7 July 2019]
 24. Woolf SH, Grol R, Hutchinson A, Eccles M, Grimshaw J. Clinical guidelines: potential benefits, limitations, and harms of clinical guidelines. *BMJ* 1999; 318: 527e30
 25. La Colla L, Albertin A, La Colla G, et al. No adjustment vs. adjustment formula as input weight for propofol target-controlled infusion in morbidly obese patients. *Eur J Anaesthesiol* 2009; 26: 362e9
 26. Jones J, Hunter D. Consensus methods for medical and health services research. *BMJ* 1995; 311: 376e80
 27. Gabriel RA, Kaye AD, Jones MR, Dutton RP, Urman RD. Practice variations in anesthetic care and its effect on clinical outcomes for primary total hip arthroplasties. *J Arthroplasty* 2016; 31: 918e22
 28. Helwani MA, Avidan MS, Ben Abdallah A, et al. Effects of regional versus general anesthesia on outcomes after total hip arthroplasty: a retrospective propensity- matched cohort study. *J Bone Jt Surg Am* 2015; 97: 186e93
 29. Aynardi M, Jacovides CL, Huang R, Mortazavi SM, Parvizi J. Risk factors for early mortality following modern total hip arthroplasty. *J Arthroplasty* 2013; 28: 517e20
 30. Brinker MR, Reuben JD, Mull JR, Cox DD, Daum WJ, Parker JR. Comparison of general and epidural anesthesia in patients undergoing primary unilateral THR. *Orthopedics* 1997; 20: 109e15
 31. Chen WH, Hung KC, Tan PH, Shi HY. Neuraxial anesthesia improves long-term survival after total joint replacement: a retrospective nationwide population- based study in Taiwan. *Can J Anaesth* 2015; 62: 369e76
 32. Harsten A, Kehlet H, Ljung P, Toksvig-Larsen S. Total intravenous general anaesthesia vs. spinal anaesthesia for total hip arthroplasty: a randomised, controlled trial. *Acta Anaesthesiol Scand* 2015; 59: 298e309
 33. Haugthorn BD, Schairer WW, Nwachukwu BU, Hellman MD, Levine BR. Does neuraxial anesthesia decrease transfusion rates following total hip arthroplasty? *J Arthroplasty* 2015; 30: 116e20
 34. Hole A, Terjesen T, Breivik H. Epidural versus general anaesthesia for total hip arthroplasty in elderly patients. *Acta Anaesthesiol Scand* 1980; 24: 279e87
 35. Khatod M, Inacio MC, Bini SA, Paxton EW. Prophylaxis against pulmonary embolism in patients undergoing total hip arthroplasty. *J Bone Jt Surg Am* 2011; 93: 1767e72
 36. Modig J, Maripuu E, Sahlstedt B. Thromboembolism following total hip replacement: a prospective investigation of 94 patients with emphasis on the efficacy of lumbar epidural anesthesia in prophylaxis. *Reg Anesth Pain Med* 1986; 11: 72e9
 37. Liang C, Wei J, Cai X, Lin W, Fan Y, Yang F. Efficacy and safety of 3 different anesthesia techniques used in total hip arthroplasty. *Med Sci Monit* 2017; 23: 3752e9
 38. Dauphin A, Raymer KE, Stanton EB, Fuller HD. Comparison of general anesthesia with and without lumbar epidural for total hip arthroplasty: effects of epidural block on hip arthroplasty. *J Clin Anesth* 1997; 9: 200e3
 39. Walton JK, Robinson RG. Some observations on urinary retention and deep hip sepsis following total hip replacement. *Aust N Z J Surg* 1982; 52: 130e3
 40. Walts LF, Kaufman RD, Moreland JR, Weiskopf M. Total hip arthroplasty. An investigation of factors related to postoperative urinary retention. *Clin Orthop Relat Res* 1985; 194: 280e2
 41. Davis FM, Laurensen VG, Gillespie WJ, Wells JE, Foate J, Newman E. Deep vein thrombosis after total hip

- replacement. A comparison between spinal and general anaesthesia. *J Bone Jt Surg Br* 1989; 71: 181e5
42. Harsten A, Kehlet H, Toksvig-Larsen S. Recovery after total intravenous general anaesthesia or spinal anaesthesia for total knee arthroplasty: a randomized trial. *Br J Anaesth* 2013; 111: 391e9
 43. Thorburn J, Loudon J, Vallance R. Spinal and general anaesthesia in total hip replacement: frequency of deep vein thrombosis. *Br J Anaesth* 1980; 52: 1117e21
 44. Macdowell AD, Robinson AH, Hill DJ, Villar RN. Is epidural anaesthesia acceptable at total hip arthroplasty? A study of the rates of urinary catheterisation. *J Bone Jt Surg Br* 2004; 86: 1115e7
 45. Williams A, Price N, Willett K. Epidural anaesthesia and urinary dysfunction: the risks in total hip replacement. *J R Soc Med* 1995; 88: 699e701
 46. Feller JA, Parkin JD, Phillips GW, Hannon PJ, Hennessy O, Huggins RM. Prophylaxis against venous thrombosis after total hip arthroplasty. *Aust N Z J Surg* 1992; 62: 606e10
 47. Haas S, Holberg G, Kreutz R, et al. The effects of timing of prophylaxis, type of anesthesia, and use of mechanical methods on outcome in major orthopedic surgery subgroups analyses from 17,701 patients in the XAMOS study. *Vasc Health Risk Manag* 2016; 12: 209e18
 48. Sarmiento A, Goswami A. Thromboembolic disease prophylaxis in total hip arthroplasty. *Clin Orthop Relat Res* 2005; 436: 138e43
 49. Dalldorf PG, Perkins FM, Totterman S, Pellegrini Jr VD. Deep venous thrombosis following total hip arthroplasty. Effects of prolonged postoperative epidural anesthesia. *J Arthroplasty* 1994; 9: 611e6
 50. Fredin H, Rosberg B. Anaesthetic techniques and thromboembolism in total hip arthroplasty. *Eur J Anaesthesiol* 1986; 3: 273e81
 51. Modig J, Hjelmstedt A, Sahlstedt B, Maripuu E. Comparative influences of epidural and general anaesthesia on deep venous thrombosis and pulmonary embolism after total hip replacement. *Acta Chir Scand* 1981; 147: 125e30
 52. Modig J, Borg T, Karlstrom G, Maripuu E, Sahlstedt B. Thromboembolism after total hip replacement: role of epidural and general anesthesia. *Anesth Analg* 1983; 62: 174e80
 53. Wille-Jorgensen P, Christensen SW, Bjerg-Nielsen A, Stadeager C, Kjaer L. Prevention of thromboembolism following elective hip surgery. The value of regional anesthesia and graded compression stockings. *Clin Orthop Relat Res* 1989; 247: 163e7
 54. Basques BA, Toy JO, Bohl DD, Golinvaux NS, Grauer JN. General compared with spinal anesthesia for total hip arthroplasty. *J Bone Jt Surg Am* 2015; 97: 455e61
 55. Brueckner S, Reinke U, Roth-Isigkeit A, Eleftheriadis S, Schmucker P, Siemens HJ. Comparison of general and spinal anesthesia and their influence on hemostatic markers in patients undergoing total hip arthroplasty. *J Clin Anesth* 2003; 15: 433e40
 56. Mitchell D, Friedman RJ, Baker 3rd JD, Cooke JE, Darcy MD, Miller 3rd MC. Prevention of thromboembolic disease following total knee arthroplasty. Epidural versus general anesthesia. *Clin Orthop Relat Res* 1991; 269: 109e12
 57. Park YB, Chae WS, Park SH, Yu JS, Lee SG, Yim SJ. Comparison of short-term complications of general and spinal anesthesia for primary unilateral total knee arthroplasty. *Knee Surg Relat Res* 2017; 29: 96e103
 58. Shi HJ, Xue XH, Wang YL, Zhang WS, Wang ZS, Yu AL. Effects of different anesthesia methods on cognitive dysfunction after hip replacement operation in elder patients. *Int J Clin Exp Med* 2015; 8: 3883e8
 59. Kita T, Maki N, Song YS, Arai F, Nakai T. Caudal epidural anesthesia administered intraoperatively provides for effective postoperative analgesia after total hip arthroplasty. *J Clin Anesth* 2007; 19: 204e8
 60. Benson M, Hartmann B, Junger A, Dietrich G, Bötter S, Hempelmann G. Causes of higher blood loss during general anesthesia compared to spinal anesthesia in total hip replacement: a retrospective analysis of data collected online. *Transfus Med Hemother* 2000; 27: 311e6
 61. Borghi B, Casati A, Iuorio S, et al. Frequency of hypotension and bradycardia during general anesthesia, epidural anesthesia, or integrated epidural-general anesthesia for total hip replacement. *J Clin Anesth* 2002; 14: 102e6
 62. Davis FM, McDermott E, Hickton C, et al. Influence of spinal and general anaesthesia on haemostasis during total hip arthroplasty. *Br J Anaesth* 1987; 59: 561e71
 63. Eroglu A, Uzunlar H, Erciyes N. Comparison of hypotensive epidural anesthesia and hypotensive total intravenous anesthesia on intraoperative blood loss during total hip replacement. *J Clin Anesth* 2005; 17: 420e5
 64. Flordal PA, Neander G. Blood loss in total hip replacement. A retrospective study. *Arch Orthop Trauma Surg* 1991; 111: 34e8
 65. Hamilton H, Jamieson J. Deep infection in total hip arthroplasty. *Can J Surg* 2008; 51: 111e7
 66. Hartmann B, Junger A, Benson M, et al. Comparison of blood loss using fluorescein flow cytometry during total hip replacement under general or spinal anesthesia. *Transfus Med Hemother* 2003; 30: 20e6
 67. Jones MJ, Piggott SE, Vaughan RS, et al. Cognitive and functional competence after anesthesia in patients aged over 60: controlled trial of general and regional anesthesia for elective hip or knee replacement. *BMJ* 1990; 300: 1683e7
 68. Keith I. Anaesthesia and blood loss in total hip replacement. *Anaesthesia* 1977; 32: 444e50
 69. Wong S, Tang H, de Steiger R. Blood management in total hip replacement: an analysis of factors associated with allogenic blood transfusion. *ANZ J Surg* 2015; 85: 461e5
 70. Jacob AK, Mantilla CB, Sviggum HP, Schroeder DR, Pagnano MW, Hebl JR. Perioperative nerve injury after total knee arthroplasty: regional anesthesia risk during a 20-year cohort study. *Anesthesiology* 2011; 114: 311e7
 71. Memtsoudis SG, Danninger T, Rasul R, et al. Inpatient falls after total knee arthroplasty: the role of anesthesia type and peripheral nerve blocks. *Anesthesiology* 2014; 120: 551e63
 72. Kendrić M, Šurbatović M, Đorčević D, Trifunović B, Jevčić J. Analgesic efficacy and safety of four different anesthesia/postoperative analgesia protocols in patients following total hip arthroplasty. *Vojnosanit Pregl* 2017; 74: 814e20
 73. Bonnet F, Harari A, Thibonnier M, Viars P. Suppression of antidiuretic hormone hypersecretion during surgery by extradural anaesthesia. *Br J Anaesth* 1982; 54: 29e36

74. Borghi B, Casati A, Iuorio S, et al. Effect of different anesthesia techniques on red blood cell endogenous recovery in hip arthroplasty. *J Clin Anesth* 2005; 17: 96e101
75. Modig J, Malmberg P, Karlstrom G. Effect of epidural versus general anaesthesia on calf blood flow. *Acta Anaesthesiol Scand* 1980; 24: 305e9
76. Modig J, Borg T, Bagge L, Saldeen T. Role of extradural and of general anaesthesia in fibrinolysis and coagulation after total hip replacement. *Br J Anaesth* 1983; 55: 625e9
77. Modig J, Karlstrom G. Intra- and post-operative blood loss and haemodynamics in total hip replacement when performed under lumbar epidural versus general anaesthesia. *Eur J Anaesthesiol* 1987; 4: 345e55
78. D'Ambrosio A, Borghi B, Damato A, D'Amato G, Antonacci D, Valeri F. Reducing perioperative blood loss in patients undergoing total hip arthroplasty. *Int J Artif Organs* 1999; 22: 47e51
79. Chin SP, Abou-Madi MN, Eurin B, Witvoet J, Montagne J. Blood loss in total hip replacement: extradural v. phenoperidine analgesia. *Br J Anaesth* 1982; 54: 491e5
80. Wulf H, Biscopig J, Beland B, Bachmann-Mennenga B, Motsch J. Ropivacaine epidural anesthesia and analgesia versus general anesthesia and intravenous patient-controlled analgesia with morphine in the perioperative management of hip replacement. Ropivacaine Hip Replacement Multicenter Study Group. *Anesth Analg* 1999; 89: 111e6
81. Williams-Russo P, Sharrock NE, Mattis S, Szatrowski TP, Charlson ME. Cognitive effects after epidural vs general anesthesia in older adults. A randomized trial. *JAMA* 1995; 274: 44e50
82. Williams-Russo P, Sharrock NE, Haas SB, et al. Randomized trial of epidural versus general anesthesia: outcomes after primary total knee replacement. *Clin Orthop Relat Res* 1996; 331: 199e208
83. Bugada D, Allegri M, Gemma M, et al. Effects of anaesthesia and analgesia on long-term outcome after total knee replacement: a prospective, observational, multicentre study. *Eur J Anaesthesiol* 2017; 34: 665e72
84. Hsieh JY, Lin HW. Less incidence of coronary artery disease in general anesthesia compared to spinal-epidural anesthesia after total knee replacement: 90-day follow-up period by a population-based dataset. *Eur J Orthop Surg Traumatol* 2015; 25: 927e32
85. Chelly JE, Greger J, Gebhard R, et al. Continuous femoral blocks improve recovery and outcome of patients undergoing total knee arthroplasty. *J Arthroplasty* 2001; 16: 436e45
86. Chu CP, Yap JC, Chen PP, Hung HH. Postoperative outcome in Chinese patients having primary total knee arthroplasty under general anaesthesia/intravenous patient-controlled analgesia compared to spinal-epidural anaesthesia/analgesia. *Hong Kong Med J* 2006; 12: 442e7
87. Liu J, Ma C, Elkassabany N, Fleisher LA, Neuman MD. Neuraxial anesthesia decreases postoperative systemic infection risk compared with general anesthesia in knee arthroplasty. *Anesth Analg* 2013; 117: 1010e6
88. Lingaraj K, Ruben M, Chan Y, Das SD. Identification of risk factors for urinary retention following total knee arthroplasty: a Singapore hospital experience. *Singapore Med J* 2007; 48: 213
- 89.
- Haas SB. Effects of epidural anesthesia on incidence of venous thromboembolism following joint replacement. *Orthopedics* 1994; 17: 18e20
90. Jorgensen LN, Rasmussen LS, Nielsen PT, Leffers A, Albrecht-Beste E. Antithrombotic efficacy of continuous extradural analgesia after knee replacement. *Br J Anaesth* 1991; 66: 8e12
91. Nielsen PT, Jorgensen LN, Albrecht-Beste E, Leffers AM, Rasmussen LS. Lower thrombosis risk with epidural blockade in knee arthroplasty. *Acta Orthop Scand* 1990; 61: 29e31
92. Ishii Y, Noguchi H, Sato J, Takayama S, Okada Y, Toyabe SI. Impact of anesthesia modality and mechanical venous thromboembolism prophylaxis on the incidence of symptomatic deep venous thrombosis after TKA. *J Clin Orthop Trauma* 2018; 9: 142e5
93. Nathan S, Aleem MA, Thiagarajan P, Das De S. The incidence of proximal deep vein thrombosis following total knee arthroplasty in an Asian population: a Doppler ultrasound study. *J Orthop Surg (Hong Kong)* 2003; 11: 184e9
94. Sharrock NE, Haas SB, Hargett MJ, Urquhart B, Insall JN, Scuderi G. Effects of epidural anesthesia on the incidence of deep-vein thrombosis after total knee arthroplasty. *J Bone Jt Surg Am* 1991; 73: 502e6
95. Zhou LY, Gu W, Liu Y, Ma ZL. Effects of inhalation anesthesia vs. total intravenous anesthesia (TIVA) vs. spinal-epidural anesthesia on deep vein thrombosis after total knee arthroplasty. *Med Sci Monit* 2018; 24: 67e75
96. Kudoh A, Takase H, Takazawa T. A comparison of anesthetic quality in propofol-spinal anesthesia and propofol-fentanyl anesthesia for total knee arthroplasty in elderly patients. *J Clin Anesth* 2004; 16: 405e10
97. Belmont Jr PJ, Goodman GP, Rodriguez M, Bader JO, Waterman BR, Schoenfeld AJ. Predictors of hospital readmission following revision total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2016; 24: 3329e38
98. Kayupov E, Okroj K, Young AC, et al. Continuous adductor canal blocks provide superior ambulation and pain control compared to epidural analgesia for primary knee arthroplasty: a randomized, controlled trial. *J Arthroplasty* 2018; 33: 1040e1044.e1
99. Napier DE, Bass SS. Postoperative benefits of intrathecal injection for patients undergoing total knee arthroplasty. *Orthop Nurs* 2007; 26: 374e8
100. Pope D, El-Othmani MM, Manning BT, Sepula M, Markwell SJ, Saleh KJ. Impact of age, gender and anesthesia modality on post-operative pain in total knee arthroplasty patients. *Iowa Orthop J* 2015; 35: 92e8
101. Lawrie CM, Ong AC, Hernandez VH, Rosas S, Post ZD, Orozco FR. Incidence and risk factors for postoperative urinary retention in total hip arthroplasty performed under spinal anesthesia. *J Arthroplasty* 2017; 32: 3748e51
102. Williams MR, McKeown A, Dexter F, et al. Efficacy outcome measures for procedural sedation clinical trials in adults: an ACTION systematic review. *Anesth Analg* 2016; 122: 152e70
103. Kehlet H. The stress response to surgery: release mechanisms and the modifying effect of pain relief. *Acta Chir Scand Suppl* 1989; 550: 22e8