

On the Consistency of an Impredicative Subsystem of Quine's NF Author(s): Marcel Crabbé Reviewed work(s): Source: The Journal of Symbolic Logic, Vol. 47, No. 1 (Mar., 1982), pp. 131-136 Published by: Association for Symbolic Logic Stable URL: <u>http://www.jstor.org/stable/2273386</u> Accessed: 10/04/2012 09:26

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at http://www.jstor.org/page/info/about/policies/terms.jsp

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Association for Symbolic Logic is collaborating with JSTOR to digitize, preserve and extend access to The Journal of Symbolic Logic.

# ON THE CONSISTENCY OF AN IMPREDICATIVE SUBSYSTEM OF QUINE'S NF

### MARCEL CRABBÉ

**Introduction.** NFP is the *predicative* fragment of NF. In this system we do not allow a set to exist if it cannot be defined without using quantifiers ranging over its type or parameters of a higher type. NFI is a less restrictive fragment located between NFP and NF.

We show that NFP is really weaker than NFI; similarly, NFI is weaker than NF. This result will be obtained in the following manner: on the one hand, we will show that NFP can be proved consistent in elementary arithmetic and that second order arithmetic is interpretable in NFI; on the other hand, we will prove the consistency of NFI in third order arithmetic, which is contained in NF.<sup>1</sup>

The paper is divided in four sections. In  $\S1$ , we define the concepts needed and collect a few results together in such a way that they will be ready for later use. In  $\S2$ , we will present a model-theoretic (quick) proof of the consistency of NFI (and thus of NFP). The proof will be chosen (it is not the quickest!) so as to motivate in a natural manner the details of the proof-theoretical version of it that will be presented in  $\S3$ .  $\S4$  will be devoted to the axiom of infinity in NFP and NFI.

## §1. Definitions, notations and preliminary results.

1. NF is the theory in the language of ZF whose axioms are extensionality and the instances of the comprehension schema:  $\exists y \forall x (x \in y \leftrightarrow \varphi)$ , where  $\varphi$  is a stratifiable formula and y is not free in  $\varphi$ . NFP and NFI are subsystems of NF obtained by restricting the comprehension axioms. In NFP there must be a stratification such that the indices associated to the bound variables in  $\varphi$  do not exceed the type of x and the indices of the free variables in  $\varphi$  do not exceed the type of y. In NFI the indices associated to the variables *bound or free* in  $\varphi$  do not exceed the index attributed to y. So NFP is a part of NFI.

For any stratifiable  $\varphi$ , we can prove in NFP that  $\exists y \forall z (z \in y \leftrightarrow \exists x (z = \{ \dots \{x\} \dots\} \land \varphi))$ , provided the singleton operation is sufficiently iterated. Therefore, if U is the axiom of union:  $\exists y \forall x (x \in y \leftrightarrow \exists v (v \in z \land x \in v)))$  we obtain

LEMMA 1.<sup>2</sup> NFP + U = NFI + U = NF.

If  $n \ge 1$ , NFP<sub>n</sub> and NFI<sub>n</sub> are the fragments of NFP and NFI, respectively,

Received December 10, 1979.

<sup>&</sup>lt;sup>1</sup>An introduction to NF and type theories can be found in [11] and [7]. Some useful and recent results about these systems are presented and much simplified in [2].

<sup>&</sup>lt;sup>2</sup>U can be restricted to the assertion that the union of a set of unit sets always exists :  $\forall x(x \subseteq USC(V) \rightarrow \exists y(x = USC(y))$ .

obtained by keeping as comprehension axioms only those that are stratifiable with the first *n* indices: 1, ..., *n*. E' is the axiom of NFP<sub>4</sub> asserting the existence of the set of sets whose intersection is not empty:  $\exists y \forall x (x \in y \leftrightarrow \exists t \forall z (z \in x \rightarrow t \in z))$ .

The next theorem is essentially due to Grishin [9] and can be checked easily from his proof:

First reduction theorem.<sup>3</sup> NFP = NFP<sub>3</sub> + E'; NFI = NFI<sub>3</sub> + E'.

2. Let *TP* and *TI* be the theories of types corresponding to NFP and NFI, respectively.<sup>4</sup> In the following, *T* will be either *TP* or *TI*. If  $n \ge 1$ ,  $T_n$  is the fragment of *T* built on the first *n* types: 1, ..., *n*.  $T_n^{\infty}$  is the theory of the infinite models of  $T_n$ , that is,  $T_n$  plus, for each *m*, the standard axiom expressing that there are at least *m* objects of type 1.

If  $\varphi$  is a formula of the language of T, then  $\varphi^+$  will be obtained from  $\varphi$  by raising all type indices by 1.  $T_4^{(3)}$  is the fragment of  $T_4$  whose comprehension axioms are those of  $T_3$  and those of the form  $\varphi^+$  whenever  $\varphi$  is an axiom of  $T_3$ . E is the axiom of  $TP_4$  resulting from E' by putting the type indices 1, 2, 3 and 4 at the right places. Amb (for "ambiguity") is the set of all the sentences (i.e. closed formulas) of the form  $\varphi \leftrightarrow \varphi^+$  formulated in the language of  $T_4$  (thus  $\varphi$  is in the language of  $T_3$ ).

SECOND REDUCTION THEOREM. NFP (resp. NFI) is consistent iff  $TP_4^{(3)}$  (resp.  $TI_4^{(3)}$ ) + E + Amb is consistent.

The proof follows from [13] and the first reduction theorem.

3. A model of a set of sentences of the language of  $T_n$  is a structure  $\langle M_1, \ldots, M_n; <_1, \ldots, <_{n-1} \rangle$  satisfying these sentences (in the appropriate sense), where the  $M_i$ 's  $(1 \le i \le n)$  are pairwise disjoint sets and for each i  $(1 \le i \le n - 1), <_i$  is a relation included in the cartesian product  $M_i \times M_{i+1}$ .

A model  $\langle M_1, M_2, <_1 \rangle$  of  $T_2$  is called *countably saturated* if:

 $M_2$  is countably infinite,

for every object a of  $M_2$  such that  $\{x \in M_1 | x < a\}$  is infinite there is a b in  $M_2$  such that  $\{x \in M_1 | x < b and x < a\}$  and  $\{x \in M_2 | x < b and x < a\}$  are both infinite.

LEMMA 2. (1) Two countably saturated models of  $T_2$  are isomorphic.

(2) Every countable model  $\mathcal{M}$  ( $\mathcal{M} = \langle M_1, \ldots, M_n, <_1, \ldots, <_{n-1} \rangle$ ) of  $T_n$  has an elementary extension such that, for each i ( $1 \leq i \leq n - 1$ ),  $\langle M_i, M_{i+1}, <_i \rangle$  is countably saturated.

The first part is proved in [8]. The second is obtained by taking a recursively saturated extension of  $\mathcal{M}$  (see [10] for refinements).

§2. The model-theoretical consistency proof. Let  $\varphi_1, \ldots, \varphi_r$  (r > 0) be a sequence of sentences of the language of  $T_3$ . We consider the  $2^r$  conjunctions  $\varphi_1^{\varepsilon_1} \wedge \cdots \wedge \varphi_r^{\varepsilon_r}$  where  $\varphi_i^{\varepsilon_i}$  is  $\varphi_i$  or  $\neg \varphi_i$  according to whether  $\varepsilon_i$  is 0 or 1. Let  $\psi_1, \ldots, \psi_k$  be all the conjunctions of this sort that are consistent with  $T_3^{\infty}$ . Amb $(\varphi_1, \ldots, \varphi_r)$  is the sentence  $(\varphi_1 \leftrightarrow \varphi_1^+) \wedge \cdots \wedge (\varphi_r \leftrightarrow \varphi_r^+)$ . We show that  $T_4^{(3)} + E + \text{Amb}(\varphi_1, \ldots, \varphi_r)$ is consistent. From this and the second reduction theorem we may then conclude by compactness that NFP and NFI are consistent.

<sup>&</sup>lt;sup>3</sup>For alternative reductions see [1] and [5].

<sup>&#</sup>x27;These systems are studied in [5].

From now on we fix  $r, k, \varphi_1, \ldots, \varphi_r, \psi_1, \ldots, \psi_k$  as described. A simple application of Lemma 2 gives

LEMMA 3. If  $\langle M_2, M_3, \langle 2 \rangle$  is a countably saturated model of  $T_2$ , then for each i  $(1 \leq i \leq k)$  there is a set  $M_{1i}$  and a relation  $\langle_{1i}$  such that  $\langle M_{1i}, M_2, M_3, \langle_{1i}, \langle_2 \rangle$  is a model of  $T_3 + \psi_i$ .  $\Box$ 

Let  $\langle M_2, M_3, <_2 \rangle$  be a countably saturated model of  $T_2$  and choose  $M_{11}, \ldots, M_{1k}, <_{11}, \ldots, <_{1k}$  as indicated in Lemma 3. If  $1 \le i \le k$ , let  $E_i$  be the set of the objects a of  $M_3$  such that

$$\langle M_{1i}, M_2, M_3, \langle 1i, \rangle \models \exists z^1 \forall v^2 (v^2 \in a \rightarrow z^1 \in v^2).$$

We then have

LEMMA 4. There is a model  $\langle M_2, M_3, M_4, <_2, <_3 \rangle$  of  $T_3$  such that:  $M_4$  is a collection of subsets of  $M_3$  and  $<_3$  is the standard  $\in$ -relation; each  $E_i$   $(1 \le i \le k)$  belongs to  $M_4$ .

**PROOF.**  $M_4$  may be taken as the power set of  $M_3$ , i.e. the collection of all subsets of  $M_3$ .

This works in the two cases. But, when  $T_3$  is the predicative theory one can proceed in a more economical way, since  $M_4$  may then be taken as the collection of the subsets *definable* from parameters belonging to  $M_2 \cup M_3 \cup \{E_1, \ldots, E_k\}$ .

LEMMA 5.  $T_4^{(3)} + E + \text{Amb}(\varphi_1, \ldots, \varphi_r)$  is consistent.

**PROOF.** Let  $\mathcal{M} = \langle M_2, M_3, M_4, <_2, <_3 \rangle$  be as in Lemma 4. There is exactly one  $\varepsilon$  such that  $\varphi_r^{\varepsilon_1} \wedge \ldots \wedge \varphi_r^{\varepsilon_r}$  is true in  $\mathcal{M}$ . But since  $\mathcal{M}$  is a model of  $T_3^{\infty}$  there is a unique i  $(1 \le i \le k)$  such that this sentence is  $\varphi_i$ . So, the structure  $\mathcal{M}'$   $(\mathcal{M}' = \langle M_{1i}, M_2, M_3, M_4, <_{1i}, <_2, <_3 \rangle)$  is a model of  $T_4^{(3)}$  satisfying  $\varphi_i$  and  $\varphi_i^+$ . Moreover,  $E_i$  belongs to  $M_4$ .  $\mathcal{M}'$  is thus a model of  $T_4^{(3)} + E + \operatorname{Amb}(\varphi_1, \ldots, \varphi_r)$ .

§3. The proof-theoretical version. We show here how to transform the consistency proof given above into one which uses proof-theoretical means only. Besides its own interest, such a demonstration will provide information about the relative powers of the considered systems.

1. More definitions. If *i* is a natural number and  $\varphi$  a formula of  $T_n$ , then  $\varphi^i$  will be the expression resulting from  $\varphi$  by replacing in it the type index 1 by 1*i* and  $T_n^{\infty i}$ will be the theory differing inessentially from  $T_n^{\infty}$  in that the type 1 is denoted in it by li.  $T_n^*$  is the union of the theories  $T_n^{\infty i}$  in the sense that the nonlogical axioms of  $T_n^*$  are just those  $\varphi^i$ 's that are nonlogical axioms of the  $T_n^{\infty i}$ 's.<sup>5</sup> In this section, *n* will always be 3 or 4, and *r*, *k*,  $\varphi_1, \ldots, \varphi_r, \varphi_1, \ldots, \varphi_k$  are fixed as in §2.  $\Psi$  is  $\varphi_1^1 \wedge \cdots \wedge \varphi_k^k$ , which is a sentence in the language of  $T_3^*$ . Note that the consistency of  $T_3^{\infty}$  and that  $k \ge 1$  are elementary provable.

2. LEMMA 6. It is elementary provable, from the definition of  $\Psi$ , that  $T_3^* + \Psi$  is consistent.

**PROOF.** Using the elementary proof of the quantifier elimination theorem for the theory of atomic boolean algebras (extended by some extra predicates), one

<sup>&</sup>lt;sup>5</sup>The type 1 is thus split into k parts. More precisely, the language of  $T_n^*$  has variables for the "types" 11, ..., 1k, 2, ..., n; equality symbols :=  $_{11}$ , ..., =  $_{1k}$ , =  $_2$ , ..., =  $_n$ ; relation symbols :  $\in _{11}$ , ...,  $\in _{1k}$ ,  $\in _2$ , ...,  $\in _{n-1}$ . Formulas are built as usual from atomic ones of the kind :  $x^{1i} = _{1i} y^{1i}$ ,  $x^{1i} \in _{1i} y^2$ ,  $x^m \in _m y^{m+1}$ , etc.

proves in arithmetic that  $TP_2^{\infty} = TI_2^{\infty}$  is a complete theory [3], [5]. On the other hand, Robinson's consistency theorem is elementary provable [15]. These proofs can be transferred to the typed theories.

LEMMA 7.  $T_4^*$  is a conservative extension of  $T_3^*$ . This is provable in first order arithmetic in the predicative case and in third order arithmetic when  $T_4^*$  is  $TI_4^*$ .

**PROOF.**  $T_4^*$  can be presented in Gentzen's style as a "second order" sequent calculus. If  $\Gamma$  and  $\Delta$  are finite sequences of formulas of the language of  $T_3^*$ , we let  $\Gamma \Vdash_1 \Delta$  mean that the sequent  $\Gamma \Vdash \Delta$  is provable in the sequent calculus (without comprehension or equality rules) adapted to the language of  $T_3^*$ . We introduce then a metalinguistic abstract  $\{x^3 \mid \varphi\}$  for each formula  $\varphi$  of  $T_4^*$ , if  $T_4^*$  is  $TI_4^*$ ; and for each  $\varphi$  containing no bound variables of type 4 if  $T_4^*$  is  $TP_4^*$ . We let  $\Gamma \Vdash_2 \Delta$  mean that  $\Gamma \Vdash \Delta$  is provable in the corresponding "second order" calculus (without equality or extensionality rules).

From the cut elimination theorem for the predicative second order logic (which is provable in first order arithmetic [15]) and the cut elimination for full (impredicative) second order logic (provable in third order arithmetic [14]), it is clear that if the formulas of  $\Gamma$  and  $\Delta$  are in  $T_3^*$ , then  $\Gamma \Vdash_2 \Delta$  entails  $\Gamma \Vdash_1 \Delta$ .

Let  $Eq_1$  be the axiom

$$\forall x^3 \forall y^3 (x^3 = y^3 \rightarrow \forall z^4 (x^3 \in z^4 \rightarrow y^3 \in z^4))$$

and Eq<sub>2</sub> the conjunction of the axioms for equality and extensionality for 4-typed objects. If  $\varphi$  is a formula of  $T_3^*$ , then  $T_4^* \vdash \varphi$  implies that  $\Gamma$ , Eq<sub>1</sub>, Eq<sub>2</sub>  $\Vdash_2 \varphi$  for a finite sequence  $\Gamma$  of axioms of  $T_3^*$ . If we replace, in a proof of such a sequent,  $x^4 = y^4$  by  $\forall z^3(z^3 \in x^4 \leftrightarrow z^3 \in y^4)$ , and if we then relativise the  $x^4$ 's to the unary relation  $R(x^4)$  defined by

$$\forall x^3 \forall y^3 (x^3 = y^3 \land x^3 \in x^4 \to y^3 \in x^4),$$

we obtain that  $\Gamma \Vdash_2 \varphi$  (see [15] for details). But then  $\Gamma \Vdash_1 \varphi$ , that is,  $T_3^* \vdash \varphi$ .<sup>6</sup>

LEMMA 8.  $T_4$  + Amb( $\varphi_1, \ldots, \varphi_r$ ) is consistent.

PROOF. A will abreviate  $\operatorname{Amb}(\varphi_1, \ldots, \varphi_r)$ . First, for every  $i \ (1 \le i \le k), \ T_4^* + \Psi \vdash \varphi_i^+ \to A^i$ . But, if  $T_4 \vdash \neg A$ , then  $T_4^* \vdash \neg A^i$ , for every  $i \ (1 \le i \le k)$ . Thus, if the lemma is not true, it follows that  $T_4^* + \Psi \vdash \neg \varphi_1^+ \land \cdots \land \neg \varphi_k^+$ . By the definition of the sequence  $\varphi_1, \ldots, \varphi_k$ , we have that  $T_3^* \vdash \varphi_1 \lor \cdots \lor \varphi_k$ . Hence,  $T_4^* \vdash \varphi_1^+ \lor \cdots \lor \varphi_k^+$ . So,  $T_4^* + \Psi$  would be inconsistent. This is clearly impossible in view of Lemmas 6 and 7.  $\Box$ 

If m is a positive natural number, we let  $PA_m$  be the system of mth order arithmetic (PA is  $PA_1$ ). CON(S) is the canonical sentence of PA expressing the consistency of S.

THEOREM 1.  $PA \vdash \text{CON(NFP)}$  and  $PA_3 \vdash \text{CON(NFI)}$ .

**PROOF.** Lemmas 6, 7 and 8 show that  $PA \vdash CON(TP_4 + Amb)$  and  $PA_3$ 

<sup>&</sup>lt;sup>6</sup>This proof can be adapted in order to give a proof-theoretical proof of the fact that ML (Quine's Mathematical Logic) is a conservative extension of NF.

 $\vdash$  CON(*TI*<sub>4</sub> + Amb). *PA*  $\vdash$  CON(*TP*<sub>4</sub> + Amb)  $\rightarrow$  CON(NFP<sub>4</sub>) and *PA*  $\vdash$  CON(*TI*<sub>4</sub> + Amb)  $\rightarrow$  CON(NFI<sub>4</sub>) follow from [4] and [6], where a finitary proof of Specker's result can be found. *PA*  $\vdash$  CON(NFP<sub>4</sub>)  $\rightarrow$  CON(NFP) and *PA*  $\vdash$  CON (NFI<sub>4</sub>)  $\rightarrow$  CON(NFI) is clear from [9].

§4. The axiom of infinity. In NFP, we can define the notion of natural number as usual:

$$Nn(x) \equiv_{def} \forall y (Ind(y) \rightarrow x \in y),$$

where

$$Ind(y) \equiv_{def} \{A\} \in y \land \forall x (x \in y \to x + 1 \in y)$$

and

$$x + 1 \equiv_{\text{def}} \{ z \mid \exists t \ (t \in z \land z \setminus \{t\} \in x \}.$$

The axiom of infinity says that V, the set of all sets, is not an element of a natural number or, alternatively, that  $\Lambda$ , the empty set, is not a natural number.

THEOREM 2. NFP  $\vdash \neg Nn(\Lambda)$ ; NFP  $\vdash \forall x (Nn(x) \rightarrow V \notin x)$ .

PROOF. If the axiom of union holds, then, by Lemma 1, Specker's famous proof of the axiom of infinity for NF [12] goes through.<sup>7</sup>

Suppose now that U is false and let X be  $\{x \mid A \neq x \land \forall y (y \in x \rightarrow \forall z(z \subseteq y \rightarrow Uz \text{ exists}))\}$ . X is predicatively defined when "Uz exists" abbreviates  $\exists y \forall x (x \in y \leftrightarrow \exists v (v \in z \land x \in v))$ . Note that  $\neg U$  is equivalent to  $\neg \forall z(z \subseteq V \rightarrow Uz \text{ exists})$ . The theorem will be proved if we can establish that NFP  $\vdash$  Ind(X).

It is easy to see that  $\{\Lambda\} \in X$ . Assume then that  $x \in X$ . This implies  $x \neq \Lambda$  and  $V \notin x$  (if not, U would hold). Also,  $x + 1 \neq \Lambda$  because if  $y' \in x$  and  $t \notin y'$ , then  $y' \cup \{t\} \in x + 1$ . Let y, z, t be such that  $y \in x + 1$ ,  $z \subseteq y$ ,  $t \in y$  and  $y \setminus \{t\} \in x$ . It remains to show that Uz exists. Two cases are possible. (1) If  $t \notin z$ ,  $z \subseteq y \setminus \{t\}$ ,  $y \setminus \{t\} \in x$ , and Uz exists. (2) If  $t \in z$ ,  $z \setminus \{t\} \subseteq y \setminus \{t\}$ ,  $y \setminus \{t\} \in x$ ,  $U(z \setminus \{t\})$  exists and, because finite union is predicatively definable,  $U(z \setminus \{t\}) \cup t = Uz$  exists.  $\Box$ 

Although the axiom of infinity is a theorem of NFP, it cannot be proved in this system that the set Nn ( $Nn = \{x \mid Nn(x)\}$ ) of natural numbers exists. This would in fact entail that PA could be interpreted in NFP and thus that the consistency of NFP could be proved in NFP.

Despite this, the axiom of infinity, being provable, ensures that nontrivial weak fragments of arithmetic are interpretable in NFP.

The situation is much different for NFI, since there the set of natural numbers exists. This fact permits us to interpret not only PA but also classical analysis. If we remark, moreover, that  $PA_3$  is interpretable in NF, then, from Theorem 1 follows

THEOREM 3. NFI  $\vdash$  CON(NFP) and NF  $\vdash$  CON(NFI).

<sup>&</sup>lt;sup>7</sup>Specker's argument needs U, in the restricted form of footnote 2, for showing that USC(SC(x)) and SC(USC(x)) are equipollent.

#### MARCHEL CRABBÉ

### REFERENCES

[3] M. BOFFA and M. CRABBÉ, Les théorèmes 3-stratifiés de NF<sub>3</sub>, Comptes Rendus de l'Académie des Sciences de Paris, vol. 280 (1975), pp. 1657–1658.

[4] M. CRABBÉ, Types ambigus, Comptes Rendus de l'Académie des Sciences de Paris, vol. 280 (1975), pp. 1–2.

[5] —, La prédicativité dans les théories élémentaires, Logique et Analyse, No. 74–75–76 (1976), pp. 255–266.

[6] —, Ambiguity and stratification, Fundamenta Mathematicae, vol. 101 (1978), pp. 11–17.

[7] A. FRAENKEL, Y. BAR-HILLEL and A. LEVY, *Foundations of set theory*, North-Holland, Amsterdam, 1973.

[8] V. N. GRISHIN, Consistency of a fragment of Quine's NF system, Soviet Mathematics Doklady, vol. 10 (1969), pp. 1387–1390.

[9] —, The method of stratification in set theory, Thesis, Academy of Science of U.S.S.R., Moscow, 1973 (Russian).

[10] U. OSWALD, Fragmente von "New-Foundations" und Typentheorie, Thesis, E.T.H., Zürich, 1976.

[11] W.V.O. QUINE, *Set theory and its logic*, Harvard University Press, Cambridge, Massachusetts, 1963.

[12] E. SPECKER, The axiom of choice in Quine's New Foundations for mathematical logic, Proceedings of the National Academy of Science of the United States of America, vol. 39(1953), pp. 972–975.

[13] —, Typical ambiguity, Logic, methodology and philosophy of science, Proceedings of the 1960 International Congress, Stanford, 1962, pp. 116–124.

[14] W. W. TAIT, A non constructive proof of Gentzen's Hauptsatz for second order predicate logic, Bulletin of the American Mathematical Society, vol. 72 (1966), pp. 980–983.

[15] G. TAKEUTI, *Proof theory*, North-Holland, Amsterdam, 1975.

UNIVERSITÉ DE LOUVAIN (U.C.L.)

INSTITUT SUPÉRIEUR DE PHILOSOPHIE LOUVAIN-LA-NEUVE, BELGIUM